

Fri. Dec. 17 ch

1

Repeat of n 93 expt with new DNA

Aim: To test new $\text{ch3A}(\text{lac})$ ~~for~~ for ligation & nick.

~~$\text{ch3A}(\text{lac})$ 391 growth DNA from Bill W. 0 $\mu\text{g}/\mu\text{l}$.
- called ch3A - checked!~~

100ml

✓ 10.9g
✓ 2.03g

Fresh Sugden's 10X R1 (+ATP)

0.9M Tris to pH 7.8

0.1M $\text{Mg}^{++}\text{Cl}_2$

with sterile d.w.

10mM DTT

15.4mg

+ 50mM ATP

32.0mg

To 10ml

FROZE as 01 10X R1

even though it has afflictions

∴ tried straight phage 3A (available at OD 108 in N6M C50) $\equiv$ ~~of this OD is~~ DNA
i.e. $\frac{50}{0.02 \times 1000}$ $\mu\text{g}/\mu\text{l}$ or 2.5 $\mu\text{g}/\mu\text{l}$

Used $\frac{1}{2}$
- but really 0.8

∴ so can try:

6 μl ($\approx$ 15 μg)

+ 54 μl sterile { 0.01M Tris
0.01M EDTA pH 8.0

Heat to 85°C for 5 min

← ? Too high for this

Heated phage
01

+ 2.7 μl 0.04M MgCl_2 (sterile)
+ 12 μl 01 10X Sugden's + ATP + DTT
+ 3.5 μl d.w.

but will allow good internal sticky ending

Miles Higashimoto $\frac{10}{6}$ 2.5 μ l to 12.5 Sugden's IX

nr2 DGR1 / 50 $\rightarrow$ ~~10~~ ~~10~~ ~~10~~ ~~10~~

R1/100 $\rightarrow$ 5 μ l to 5 μ l in ERI buffer

R1/200 $\rightarrow$ 5 μ l 5 μ l

R1/400 $\rightarrow$ 5 μ l 5 μ l

PL 2 Hin III Undiluted

03
DNA

03 DNA

Miles Higashimoto

DG RI

PL Hin
Lot 2

To 37°
3.35

Hydrolysed or boiled membrane

against unsterilized 0.01M Tris 0.001M EDTA pH 8

from 2.15 pm to 2.30 pm (15 min) then taken up with 50 μ l
sterile 0.01M Tris 0.001M EDTA pH 8

Heated at 85°C for 5 mins.

→ 60 μ l to 3 μ l 0.04M $MgCl_2$ sterile

12. $10 \times \text{Sugars (0.1)} + \text{ATP} + \text{DTT}$

45 μ l sterile L.W.

[chief problem is if ligation comes over in the time]

03
DNA

	RI	RI + Hin	Hin	RI + Hig	RI + Hin + Hig	Hin + Hig	RI/ 100 + Hig	RI/ 200 + Hig	RI/ 400 + Hig
3 DNA	12	12	12	12	12	12	12	12	12
Miles Higax $\frac{10}{6}$	—	—	—	2	2	2	2	2	2
DG RI/50	2	2	—	2	2	—	2($\frac{1}{2}$)	2($\frac{1}{4}$)	2($\frac{1}{8}$)
Ph Hin + III lot 2	—	2	2	—	2	2	—	—	—

1037° at
3.35 pm

Base = 6.000000


Residue as control

Stop with 3 pl

1.3 M NaOH / 0.35 M EDTA

1.3 ml
2 ml

0.7 ml
1 M



On 5 pm Off 10 am

R1

R1 + Hin

Hin

R1

+ lig

R1 + Hin + lig

Hin + lig

R1 ($\frac{1}{2}$) + ligR1 ($\frac{1}{4}$) + ligR1 ($\frac{1}{8}$) + lig

Control (no enzymes)

0.75% Agarose

HX4

~0.75 V/cm

long gel

Not bad for a first try.

Conclusions: Protein ~~gets~~ gets in the way of e.p. Control looks to be in trouble so there may be nucleases in the $CSCl_2$ which can survive $85^\circ C$ (!). No sign of ligation. All enzymes seem borderline "incomplete" e.g. Hin looks incomplete $\Rightarrow$ R1/2 didn't work in this series.

∴ for this work, I need a micro DNA method.


Note: Bill W. made a regular prep. of DNA with this phage prep $\Rightarrow$ stated that "the phenol ate it all up" - so it has nucleases in it.

∴ go over to a hot prep method. Perhaps best is try this a bit more - it is promising. If ligation is needed a storage stage after heating is needed to allow sticky ends to anneal e.g. 20

Sat. Dec. 18th

Further 3A-lac

3A-lac is: —



∴ EcoRI to completion plus any type of HindIII that gives preference of left site is a big ^{loss} help (e.g. 30% survival of right site with 100% of left would give a 30% efficient vector with no background phages).

∴ try 3A-lac DNA for differential effects of salt on HindIII or level of HindIII.


+ATP+DTT For 30 min then

First try Sugden's buffer + DFR1/50 + 2-fold dilutions of HindIII, 8 titers at usual Hind level with plus 0.1, 0.2, 0.5, 1.0 M NaCl.
 (2%) 5% 10% 25% ^{vol} 5 M NaCl added relative vol. start


Mon. Dec 20th

Further thoughts

CH3A-lac




Hi temp. anneal oligate — can use as



Trouble is background from self closure of d.

∴ better use: —




Tail with G (arc) after R1, Fill, +t in
high Mg^{++} (to avoid nick) + use

this to clone PST (or any flesh or 3' cell
enzyme product also tailed with (or G)
in presence of high Mg^{++} .

Result will be no self closures.

Still earnest is :-



→ to straight R1 cloning → pick
only whites (on gel - strain)

∴ return to characterizing DNA's &
improving tailing → get the method
of gel hybridization working.

hesley is making DNA - so this can wait.

∴ text :-

Embryonic DNA ↑ conc. plus λ visible
→ then λ at trace level for nick translated
 ^{32}P selection.

We ~~can expect to put~~ can expect to put
~ 0.4 $\mu g / mm^2$ according to Ketner & Kelly
or 0.7 $\mu g / mm^2$ according to Southern
or 5 $\mu g / mm^2$ according to our own work
i.e. up to 75 μg on our 7mm x 2mm slots
or 5 $\mu g / \mu l$.

I have DNA at 274 $\mu g / ml$ which can
be used for test, & Ch3A+lac at 650 $\mu g / ml$
→ set up up Eco RI digest of genomic & ditto of
λ → mix in appropriate proportions for test.

Mon Dec. 20th

Hybridization tests

To 37° at 11.15 a.m. ↓ 2.15 pm

$$\left\{ \begin{array}{l} 200 \mu\text{l} \left(\begin{array}{l} \text{274 } \mu\text{g/ml} \\ \approx 55 \mu\text{g DNA} \end{array} \right) \text{ } \lambda\text{B DNA} \\ + 23 \mu\text{l } 10\times\text{R1 buffer} + \sim 1 \mu\text{l } \text{DGR1} \end{array} \right\} \begin{array}{l} 0.13 \\ \text{Gen R1} \end{array}$$

$$\left\{ \begin{array}{l} 20 \mu\text{l } \text{ADA19 ch3A(+lac) DNA} (\approx 14 \mu\text{g DNA}) \\ + 2.3 \mu\text{l } 10\times\text{R1} + \sim 1 \mu\text{l } \text{DGR1}/10 \end{array} \right\} \begin{array}{l} 0.13 \\ \lambda\text{R1} \end{array}$$

Stopped with 5 min at 85°C.


For test use λ Gen R1 diluted

1x (undiluted) $\rightarrow \frac{1}{10} \times$ (with ~~10~~ DNA buffer)

Letto Control (no RT)

Add 0.13 λR1 at $1/1000$, $1/10,000$, $1/100,000$ level diluted in Gen R1.

10 μl Gen R1 to ~~100 μl Gen R1~~ ^{100 μl Gen R1} DNA buffer




SS. BA 85 for transfer - store dry
 - works to 10^{-12-14} pg DNA in reconstruction
 - ~ 4mm gel - horizontal with agarose 65°
 (203-05) insertion. with agar seal.

1.5M NaCl } solⁿ I 30-45 min with 0.2%
 0.5M NaOH } ~60 hr 1%

3M NaCl } solⁿ II (critical to give
 0.5M Tris pH7 } Eganwapor multipar
 gel v. brittle) slow shake with
 ~2 hr. 0.2% no visible NaOH
 ~6 hr. 1% coming out
 overnight 2%

Ndrosall. L.W. 1-2 min. then to 10xSSC just
 before use (a few min.)



- Gel on 2 wicks (wet)
- Pour 10x SSC on top - no bubbles under gel
- Ndrosall. - no bubbles
- 2 x top 3mm - no bubbles
- ~2 paper towels x20
 As stack + ~1kg 10x10

10^5 to 2×10^5 m.w. transfer time 10-14 hrs.
 20×10^6 15-20 hrs.

Overnight ok. for most papers.
 Sometime top stack replaced for 6-7 hr

e) Remove papers & allow (on 3mm) multiple to dry at room temp.

f) Vac. dehydrate, & heat at 80°C for 2 hrs.
 with lib scales to seal BSA
 into filter.

Jerry uses PM mix: - 3x SSC 0.02% BSA 0.02% Ficoll 400
 0.02% PVP 360

- like better than 40% formamide

- purify probe on G50 or on gels - crude rx. mix gives high background.

Typically $1-10 \times 10^5$ cpm. / 1-2 ml
 12 hr

Can have 0.1-0.5% SDS in final hybrid? mix as safeguard.

Botchan uses 68°C 6x SSC 0.5% SDS 0.001M EDTA
 0.02% Ficoll 0.02% BSA 0.02% PVP



make up & sterilize 100 ml.

60 ml 10x SSC 0.1 ml 1M EDTA
 2.5 ml 20% SDS 20 mg Ficoll F4375
 ~ 37.4 ml liq. 20 mg BSA A-H303
 20 mg PVP 360

↑
 Add last


Signum
 from
 Jerry S.

0.1% hybridization
 cd.



10/22/01
10/22/01
10/22/01

10/22/01



The 1/10,000 contains $\sim 10^{-9}$ g λ DNA
 per slot.
 Gel shows n13 DNA is good, R1 acts as it well, all is OK.
 11.0am to solution I 1.5M NaCl 0.5M NaOH

11.30am to sol. II 3M NaCl 0.5M Tris pH7

3.0pm transferred to 10xSSC transfer system

10.0am Millipore off to 80%
 Gel shows nothing until returned
 then high MW's are visible.

11.30am Millipore to 80°C over Ariele in vacuo

30
5
6
7
8
9
10

Tube on giger

✓ 0.01 mR/hr

0.55

34

29

1.8

0.5

0.7 mR recovered

} Pooled as [10 21 7 Probe (32P)]

Added 7 μ l of 10% DEPC in EtOH.

4000 cpm / μ l

(Bray's
DeMumbr)

5 μ l 20091

Total incorp. $\therefore \sim 1 \mu$ Ci or 10%.

Chen

$\sim 60 \mu$ l to Jeff Ross

Tues Dec 21st

Remake of nick translated probe

Take 2.5 μ Ci of each dNTP

Dec. 8th dATP is 1 μ Ci/ μ l $\therefore$ took 5 μ l of each
 dCTP is "
 dGTP is "
 dTTP is "

Blow down

HF 9-133

Nick translation buffer ~~10X~~ 10X (0.5M Tris 0.05M
 $MgCl_2$ + BSA) in 100 μ l portions. Add 0.7 μ l ME &
 dilute to 1 ml.

✓ 1 μ l Charon ~~3A~~ 3A DNA (0.7 μ g)
 ✓ to 100 μ l 1X Nick translation
 ✓ 2 μ l DNA pol I } Lot 3443 P-L (DNA pol.
 m151 } taken up to 10 μ l E. coli B)

1 pm to 37°



↑
 mistake

to 1.50 pm

g 100 Sephadexed

to 0.3M NaCl
 0.01M Tris
 0.001M EDTA
 pH 7






Wed. Dec. 22nd

Characterization of probe

- ✓ 2 μ l 10 $\times$ BAM buffer (m93)
 - ✓ 3 μ l ADA 19 d3A DNA ($\approx 2 \mu$ g DNA)
 - ✓ 5 μ l σ 21 Probe (20,000 cpm)
 - ✓ 8 μ l ~~BAM~~ d.w.
 - ✓ 2 μ l BAM ~~(m93)~~ H-1 (n7)
- 1.15 pm to 37°C - 2.30 pm



Off 6.45 pm
Fast dried & set to
expose at 6.45 pm - 8 pm
8 pm $\rightarrow$ 8 am

Recalled in Cat Br & photo!

O.K. to use - but clearly 15°C w'd
be better.

Fri. Christmas Eve

Bag + ~ 30 ml $2 \times \text{SSC} / \frac{1}{2} \% \text{ SDS}$. Drained.
 Readded. Drained. Floated on & turned over in
 $2 \times \text{SSC} / \frac{1}{2} \% \text{ SDS}$. Incubated $\sim \frac{1}{2} - \frac{3}{4}$ hrs. Ditto
 at 70°C . Drained (bag sealed) & replaced.
 Super⁺ was colder than filter so counts
 that are left on for 2nd wash are hard
 to. After second $\geq \frac{1}{2}$ hr 70° wash rinsed
 for a few mins. in $2 \times \text{SSC}$ & assembled
 on 3MM for drying. Did not hold together, so
 stapled between two papers. Rough autorad.
 from ~ 3 pm to 10 am. through 3MM. (dd film)

[See 031 for results]

Back on for more exposure without
 paper between $\sim 11:30$ am $\rightarrow$ 9 am
 25° 27°
 (new film)

12:15 pm back to hybridⁿ with ~ 0.25 ml more of buffer.

Bill Sugden's tips

$\sim 10^6 - 10^8$ per sheet. (~~0.2~~ 1 μ g)

baked
Wet filter with 6 x SSC
Put in bag with 150ml hybrid. solⁿ
at 70°C ~ 12 hr (or overnight)
Full size + 3-5 ml hybrid. mix
Overnight or 24 hr.

Remove 32 P

Wash 1% SDS 2 x SSC
— many times every 15 mins.
for ~ 6 hr.

Took the baked gel transfer Millipore
(broken badly by vac. release !!)

~ 50 pm transferred to 150ml Denhardt's BSA, PVP buffer + to $65^\circ\text{C} - 70^\circ$

9.30 am removed buffer & replaced with
3 ml hybridⁿ (i.e. + SDS) Denhardt
+ 200 μ l probe 0.2 i.e. 0.8×10^6 c.p.m.

Mistake 1 — should have denatured
the probe first !!

11.50 am i.e. took 200 μ l more of probe + 22 μ l 2M NaOH

11.56 am — then plus 4 μ l 1M HCl ~~1M Na₂HPO₄~~
1M Na₂HPO₄ 1M NaH₂PO₄

Thurs. Dec. 23rd

Check of new DNA's & of enzymes

Following enzymes are needed: -

(R1)

DGR1 is good & 01 Sugden's
10X + ATP + DTT is checked.

Ligase 10

checked extensively & it's the
best we have

DNA Polymerase I

not yet checked { Behring

Bob Wells t.t.

tested well ~ n 109

Harvey will check this.

Check new DNA's with R1 / ⁺ligase conditions
plus no enzy. control

Conditions are ~~for~~ each DNA

~ 2 µg DNA

2 µl 10X Sugden's R1 + ATP + DTT (01) 9 µl d.w + 1 µl

2 µl ligase 10 / 6 [2 µl ligase 10 + 10X R1]



2 µl DGR1 / 50 (n 92)

Water to 20 µl

ligase 10/
69pl d.w. 1pl DRI+0.1
2pl ligase 1028 DRI/
50~~28~~

DRI

1pl

E. coli preserving
buffer 50pl

XR1+01

Fri. Dec 24th

DNA tests

evening
50 μ l

3A-lac

Hb4

Hb9

Hb11

1.35 μ g/ μ l - take 21.075 μ g/ μ l - take 2350 μ g/ μ l - take 6425 μ g/ μ l - take 5

- these were made on the 20th & so have not
had v. long to sticky end in the cold.

	3A-lac	Hb4	Hb9	Hb11
10XRI	2 ✓	2 ✓	2 ✓	2 ✓
DNA	2 ✓	2 ✓	6 ✓	5 ✓
d.w.	12 ✓	12 ✓	8 ✓	9 ✓
ligase 10/6	2 ✓	2 ✓	2 ✓	2 ✓
DGR 1/50	2 ✓	2 ✓	2 ✓	2 ✓

To 37° at 10.30am
1
11.40am to ice

stopped at 11.50am with 2 μ l 3M NaOH
made up in 1mM BTA

On 12 noon Off 2.30pm

3A-lac

Hb4

Hb9

Hb11

0.75% Agarose

Hx4

3V/cm

Re-soaked
in Hx4
start of
up again
3.30pm at
1/2 V/cm ✓

Conclusions: ligation is good in the 3 Hb clones

& separate in 3A-lac. No R₁ ~~separate~~ sites in Hb

①

Unlabeled
genomic DNA

DNA/10

$\frac{1}{4}$ $\frac{2}{100}$ $\frac{2}{1000}$ $\frac{2}{10,000}$ $\frac{2}{1,000,000}$ $\frac{2}{10,000,000}$ $\frac{2}{1,000,000,000}$ $\frac{2}{10,000,000,000}$ $\frac{2}{1,000,000,000,000}$

③

②

①

Unlabeled genomic DNA

DNA/10

1/1
1/2
1/3
1/4
1/5
1/6
1/7
1/8
1/9
1/10
1/11
1/12
1/13
1/14
1/15
1/16
1/17
1/18
1/19
1/20
1/21
1/22
1/23
1/24
1/25
1/26
1/27
1/28
1/29
1/30
1/31
1/32
1/33
1/34
1/35
1/36
1/37
1/38
1/39
1/40
1/41
1/42
1/43
1/44
1/45
1/46
1/47
1/48
1/49
1/50
1/51
1/52
1/53
1/54
1/55
1/56
1/57
1/58
1/59
1/60
1/61
1/62
1/63
1/64
1/65
1/66
1/67
1/68
1/69
1/70
1/71
1/72
1/73
1/74
1/75
1/76
1/77
1/78
1/79
1/80
1/81
1/82
1/83
1/84
1/85
1/86
1/87
1/88
1/89
1/90
1/91
1/92
1/93
1/94
1/95
1/96
1/97
1/98
1/99
1/100

②

Sat. Dec. 25th (Christmas Day!)

Transfertext

Results show even on first try, reasonably good transfer & sensitivity. No background - so very prolonged washing is not needed.

- Total washes were
- i) overnight prewash
 - ii) Several rather crude rinses with $\frac{1}{2}$ $\times$ $\frac{9}{10}$ SDS 2xSSC
 - iii) Transfer to floating on Nitro
 - iv) One $\sim \frac{3}{4}$ hr 70°C wash - probably this one was all that was needed
 - v) Second ~ 1 hr 70°C wash Nitro
 - vi) Brief 2xSSC room temp wash.

However it is clear that the probes need to be at least 1 order of magnitude better, or more DNA. Can just see 1 copy in 10^4 .
Did not overload gel with $\sim 4 \mu\text{g}$ in a slot but this was only $\sim 0.3 \mu\text{g}/\mu\text{l}$ & can expect to go to 5 μg .

i.e. in next test use ppt^d R1 DNA at higher concⁿ.
Fri 3pm $\rightarrow$ (1) $\checkmark$ use ppt^d R1 DNA at higher concⁿ.
Exposed Sat am $\rightarrow$ (2) $\checkmark$ New film $\rightarrow$ Marasm. $\rightarrow$ (3) $\rightarrow$ Marasm

Mon. Dec. 27th

Euk DNA cells

In view of X hybrid⁺ Southern test success decided to try mouse DNA for cDNA probe. (Isolate: -

~ 5g mouse liver (~5mg DNA according to DeMars)

Grind in liquid N₂.

Add to { 0.5M EDTA } pH 7 at 60-70-80°
50 ~~ml~~ ml { 2% SDS } (do not exceed 80°)

Decanted to ~5g equiv^s of liver which is fairly reasonable ~~pld~~ mess.
Held at 80° - 70° for ~30min

RAF
RNA'se m13 (reheated) added to 10µg/ml
i.e. 500µl of ~~100~~ 1µg/µl

- allowed to cool to ~60°C

added PRONASE m ~~141~~ I41 to 50µg/ml i.e.

~12.40pm

500µl of 5µg/µl - no effect - must be too low (it is 4x less than I used previously)

sh'd try 20 ~~100~~ µg/µl So added to 50ml 1mg solid

- still no sign of activity. Put to 50° bath. 1.10pm

Try Stafford's conditions. - no sign of digestion.

Try Stafford's conditions.

Stafford's procedure: -

Make 0.5% Sarkosyl in 0.05M EDTA pH 8
 Added Proteinase K to 100 μ g/ml. (5 ng to 50 ml)
 Heated to 50°C & added powder
 slowly. Temp. fluctuated rather a bit
 & rose to 60°C - but all was added & went
 very viscous. ≥ 5 g / 50 ml. Added ~ 1 mg extra proteinase
 K & left from 4.0 pm to ~ 4.30 pm at
 $\sim 50^\circ$. Still chunky & η reduced a lot, but
 there will be glycogen present etc.

Better procedure is probably a compromise
 - {0.1M EDTA pH 8} - 70°C until well
 {Sarkosyl
 dispersed} $\downarrow$ to 50°C for proteinase K
 addition & digestion.

By am, the proteinase K looked
 good. Proteinase didn't go & wouldn't
 get viscous even with Sarkosyl
 - aban for.

Tues. Dec. 28th

DNA (contd)

Cent'd Stafford mix on IEC to remove (small) amount of unhydrolyzed. Ferritin? & glycogen? did not come down. η still same as yesterday.

Extracted ~~3X~~ with aq. sat'd (0.1M Tris pH 8 & a bit of Tris-HCl) in 0.1% HO Quinidine phenol. No viscosity. Try again.

$\sim 5.5g$ powdered & to 0.1M EDTA pH 8 mag stored (max) with $\downarrow$ 0.5% Sarkosyl at $65^\circ - 75^\circ C$. Took ~ 20 min. to add all powder to 50ml. Held 15 min at $\sim 70^\circ$ - ~~it went to a ppt.~~ not the gooey mass of first stage. I suspect too low Sarkosyl. Temp. to $55^\circ C$ then added 5mg proteinase K.

It was! I added 0.5% of 30% 

So η re-added. η immediately began to $\uparrow$. Repeat at 50° with 1% Sarkosyl & with enzyme.

Trial 3

0.1M EDTA pH 8

1% Sarkosyl

(made from 30%)

Proteinase K

100 µg/ml

i.e. 3.3%

Temp 50°-55°C

Slow powder addition.

2.35

|

max. mag. stir.

Very much better / strings up to 4 feet!


3.05

Then to 55° incubation

3.15 pm - 5.0 pm

3 x Phenol pouring off DNA (extremely viscous!)
 Cent. is left with last phenol in old decanted & recent!

Post dialysis



Conc.

$$0.25 \times 50 \times 100 \mu\text{g/ml}$$

or 1.25 mg/ml

but this will be
over-estimate as dialysis
was not extensive

Pre-dialysis

O.D. 10 μ l in 1000 μ l

310	.04
300	.08
290	.17
280	.38
270	.51
260	.54
250	.52
240	.67

0.41 MOW DNA
dialysed

Wed Dec 29 uk

DNA (contd.)

24:1 CHCl_3 / Isobutanol $\sim$ equal vol.
extⁿ gave an interface in which some
objectionable pink was removed but no
DNA. Decanting was again easy (uphill still!)
Each time more came down. ? centⁿ time
On last (3rd) extⁿ centⁿ 10,000 rpm on
Sorvall for 15 min. No improvement
pH on paper at this point is 9.0 - watch
that keep phenol at 7.5 in future.

Vac'd off CHCl_3 / Isobutylalc. without
problem. ~~as~~

$\sim$ 40 ml - 50 ml DNA $\therefore$ added 0.5 ml $\sim$ 13 RNA^{ase}
(1000 $\mu\text{g}/\text{ml}$) to give $\sim$ 10 $\mu\text{g}/\text{ml}$. ($\frac{1}{10}$ of Stafford)
10.30am to 37°

12.30pm

Phenol extracted 2x more
 CHCl_3 / Isobutanol 3x
Vac'd. [then must shear or R1. Need a
check on this]

Recovered 40 ml. Added 400 μl 10% DEPC
 $\rightarrow$ to cold as

[0.41 Mouse DNA]

The length of string has dropped substantially
since RNA^{ase} - now $\sim$ 2 inches.

4.30pm

$\sim$ 5 ml in to freeze in cold v. 4-l. $\sim$ DNA buffer

32 850 cpm / 10 μ l

$\therefore$ $\left(\frac{3,285}{\mu\text{l}} \right)$ $\left[0.43 \text{ } ^{32}\text{P} \right]$
 Total 3×10^6 ($\sim 10\%$)

Thurs Dec 30th

$\rightarrow$ 100 μ l dial^d from 10.25 am $\rightarrow$
 against DNA buffer $\left. \begin{array}{l} .01 \text{ M NaCl} \\ .01 \text{ M Tris} \\ .001 \text{ M EDTA} \end{array} \right\} \text{pH } 8$

Recovered ^{none} ~~nothing~~ (got out off ring)
~~blew down to~~ ~~scrapped~~ - except for one
 as λ control at 8.45

$\rightarrow$ 200 μ l + 600 μ l EtOH + cooled (salt comes out) - back up + 20 μ l d.w.

Dialyzed v. d.w. from 11.25 am to 11.55 am

Recovered 17 μ l. at **23,000 cpm / μ l**

2 μ l
 45,800 cpm
 in Bragg

Future probes - make more
 concd. - also dialyze
 to 10^{-3} M EDTA alone with
 smaller column & fractions

0.43 (μP)
 Concd.

Wed. Dec. 29th

New nick translated probe

Need good probe for assaying RT completion of euk. DNA.

Dec. 8th 1 pCi / μ l $\therefore$ now $3.5 \mu\text{l} \times 2.5 / 2.5 \mu\text{Ci}$
 $\sim 10 \mu\text{l}$

$\therefore$ 10 μl of each dNTP listed below
 (i.e. $4 \times 0.35 \times 10 = 14 \mu\text{Ci total}$)

H# 9-133 10X nick trans. buffer 100 μl ✓
 + 900 μl d.w. + 0.7 μl ME ✓

Don't ten
 p next
 time

100 μl ✓ ✓

1 μl Chason 3A (ADA 19) $\sim 0.7 \mu\text{g}$

2 μl m151 DNA polymerase

11.50am

To $15^\circ\text{C} \pm 2^\circ\text{C}$

12.35pm

DEPC to 0.1%

0.3M NaCl 0.01M Tris 0.001M EDTA pH 7

Collected in ml then DEPC to 0.1% again

3	< .02 ml	6	15	2
4	< .02	7	35	
5	0.55	8	7	

Vol 920 μl + 10 μl 10% DEPC

32P
 1.45 g
 50-1000
 windows

Thurs. Dec. 30th

Eco R1 digestion tests on mouse DNA

DNA is 1.25 $\mu\text{g}/\mu\text{l}$ (still $\frac{1}{4}$ of max^m permissible) - but check: -

~~Control~~

Dialysed control (no eng.)


2 1/20

~~Undigested~~

Undigested (eng. + λ)

Dialysed bulk (eng. but no $^{32}\text{P}\lambda$)

Dialysed test (eng. + λ).



DNA

16



CHINA - THE NATIONAL
ANTHROPOLOGICAL ARCHIVES
1000 UNIVERSITY AVENUE
WASHINGTON, D.C. 20004
TEL: 202-619-1300
FAX: 202-619-1301

CHINA - THE NATIONAL
ANTHROPOLOGICAL ARCHIVES
1000 UNIVERSITY AVENUE
WASHINGTON, D.C. 20004
TEL: 202-619-1300
FAX: 202-619-1301



CHINA - THE NATIONAL
ANTHROPOLOGICAL ARCHIVES
1000 UNIVERSITY AVENUE
WASHINGTON, D.C. 20004
TEL: 202-619-1300
FAX: 202-619-1301

On 2.10 pm off Early am

 λ control (0.43) (^{32}P) λ R1 Digest (^{32}P)

46,000 cpm

0.75% agarose

Hx4

{ Mouse DNA + λ (^{32}P) } R1 Digest

46,000 cpm

1 v/cm

Mouse bulk R1 Digest

Control mouse DNA (no eng.) /20

Control mouse DNA (no eng.) 1x Exposed Mar Jan 3

9am $\rightarrow$

On 2.05 pm off 4.35 pm

Smaller slots

 λ R1 Digest ^{32}P

(residues)

0.75% agarose

Hx4

{ Mouse DNA + λ } R1 Digest ^{32}P

Bulk Mouse DNA

4 v/cm

Control Mouse /20 no eng.

Brief stain.

Control Mouse /1x no eng.


Fast dry.

Autorad - ~6 pm

~8 am

Conclusion:


Digestion goes ok. There are quite a lot of RNA fragments in this material so DNA conc. is probably closer to 0.5 $\mu g/\mu l$. The starting DNA looks ok.



→ 3pm ~10min in 2xSSC 3.25 Jret at 80 under vac

3hr is not enough. Use overnight

~5.30pm




Thurs. Dec. 20th

DNA transfer tests (mouse)

0.45 { Bulk Digest } 200 μ l + 2 μ l { 0.13 γ R1 digest }

On 255 pm Off 9.30a



After transfer hybridize to both probes. Good control needed for hybridization - but in future can add native DNA to get control for digestion also.

10.30 $\rightarrow$ 10.55

Solution I

11.0 $\rightarrow$ 11.25

Solution II

3 $\frac{3}{4}$ millipore (had some finger prints on it)
3 $\frac{1}{4}$ " x 3 $\frac{1}{4}$ " Top paper
3 $\frac{1}{4}$ " x 3 $\frac{1}{4}$ " Bottom paper
2 x 1 lb 200 g Paper towels

12 noon

Conclusion
looks like a possible positive

Allowing 30% for DNA content

20 pl of $\phi 51$ 10x R1 Mouse $\approx$ 200 μ g

if 100%

60 μ g

Fresh 0.7% Agarose made

1.50 pm off 10.45 am

Use for
gDNA

1/100 $\phi 137$ R1

$\phi 51$ 10x R1 Mouse (20 pl)

$\phi 51$ 5x R1 Mouse (20 pl)

$\phi 45$ R1 Mouse (20 pl)

$\phi 45$ R1 Mouse (20 pl + 1 pl of 1/100 $\phi 137$ R1)

$\phi 51$ 10x R1 Mouse (20 pl + 1 pl of 1/100 $\phi 137$ R1)

1/100 $\phi 137$ R1

0.7% Agarose
Hx4
0.5 v/cm

Use for
a DNA

Conclusion
Two 10x loads is
a little too much

Fri. Dec. 31st

New Year's Eve

Mouse R1 transfer.

Gels show no problem with over load, $\therefore$ $\uparrow$ conc.

700 μ l $\approx$ 45 ^{Bulk} Mouse R1 + 80 μ l 3M NaAc pH 8 $\rightarrow$ to -20°C
+ 2.4 μ l ^{alcohol}

$\sim$ 2 hrs. later cent. off at several top.


After decanting & tip wiping, took up ^{with} 70 μ l ~~buffer~~ (0.01M Tris 0.001M EDTA sterile). Cent. off glycogen successfully at top speed.

OD. 2 μ l in 1000 μ l DNA buffer } }

(OD alcohol super^t 10 μ l in 1000 μ l DNA buffer OD $< .02$)

310	.025
300	.045
290	.125
280	.45
270	1.0
260	1.2 (264 max)
250	.79
240	.35
230	.25
220	.80


This OD is almost 2x expected value.



51 10X
R1 Mouse

DNA


52



///

looks as if 10X is overexposed.

///



V. effective transfer

Sat. Jan 1st 1977

DNA transfer

10.00am 11.10 gel stained

11.15am → 11.45am

11.45am → 12.15pm

Solⁿ I - { 1.5M NaCl

Solⁿ II - { 0.5M NaOH

{ 3M NaCl

{ 0.5M Tris pH 7

Glass Plate

Bottom 3MM

Top 3MM

Multipore

Papertowels




$3\frac{1}{4} \times 12$ " (i.e. slide width)


$3\frac{1}{4} \times 4$ " (i.e. slide size)

$4\frac{1}{4} \times 4\frac{1}{4}$ "

4" x width




Set up by 12.25pm 2 lbs.



11.00am Off → to 2xSSC. [All 20x SSC was through].

11.30am to 80° under vac (plus older ϕ (9 test) reheat)

2.0pm off Cut




Also from
 $\phi 49$

2.30pm to 70°C 9.15am


DNA

Result shows excellent transfer
& successful hybrid? E.p. is good on
the 1X mouse R1 but is doubtful
on the 10X. Too early to tell - but proceed
with other



Results back with gels

Mon. Jan 2nd



Expelled wash
+ sealed 95%. Closed
off with hemostats

(+N)
φ53 pieces
+ φ49

9.48am 100 μ l φ43 (32 P) γ probe (300,000 cpm)
+ 15 μ l 2M NaOH

9.53am plus 30 μ l φ25 neutralizing buffer { 1M HCl final
2M Phos pH ~ 8
Then plus 2ml of φ17 hybridization buffer }

Transferred to bag. Too little vol. to separate
papers - would be okay with one.

Added 1ml more + tried again. Not good
but left - since this is a trial.

10.30am
to 70°

9.0am off - Washed in the bag with ~30ml 2xSSC
~30ml 0.5%
~10ml SDS

10.15am to 70° with ~30ml 2xSSC + 0.5% SDS

↓
11.45am

12.0pm to 70° with ~40ml.
(temp. a bit down)

↓
2.30pm

Rinsed in 2xSSC. Dried & to autovar 3.10pm
Wed 8.30am → Sat 10 → N. week

8.10am
↑

→ 3.45pm Fri. Took off. Washed 2x 2xSSC + $\frac{1}{2}$ %
set up 3.45pm at 70°C. Second wash
4.15pm — 5.15pm 2xSSC rinsed. Dried
to autoclave. 5.30pm Fri → 10am Sat → 1 week
(see 050)

Thurs. Jan 6th

Real test

12 noon

Rest of globin strip to Denhart's
at 70°C ~ 100 ml.

8:30 am

Drained & set up

Hemostat
with rubber
cover.



1.8 ml of (32 P) cDNA (ss) globin from Harv
HF 10-54 2.2×10^6 in ~~100~~ 0.001 M EDTA
sep 27

$\therefore$ make up } 12XSSC + 1% SDS + Ficoll + BSA + PVP
50 ~~100~~ ml } .02% .02% .02%

2xHybⁱⁿ
buffer 12XSSC

17.5 ml d.w.
30 ml 20X, 90mg F, 90mg BSA, 90mg PVP
Sterilized & added 2.5 ml 20% SDS

10:0 am

Probe + equal vol 2xHybⁱⁿ buffer
& set up at 70°

4:30 am

Web. off - extruded hybridⁱⁿ mix $\uparrow$ - sealed
& re-used - for $\phi 49$ piece

~9:15 am

$\phi 49$ piece to 6XSSC Denhart's at 70° (wash for
1 hr only - $\sim 10:15$ am to old hybridⁱⁿ from 6 to 70°
rough. Technique not good

Rinsed strip 2x in 2XSSC $\frac{1}{2}$ % SDS (dropped it once!)

10 am

to 70°C for 1st wash (it has a hot spot) 10:45 am 2nd wash

~2:0 pm

off & dried after 2XSSC rinses. Autorad. 2:40 pm $\rightarrow$ 10 am

DNA

OLIVER
~~Cesley~~

98

December 1976
Memo NAR-64

Specific excision of the inserted DNA segment from hybrid plasmids constructed by the poly(dA)-poly(dT) method.

H. Hofstetter, A. Schamböck, J. van den Berg and C. Weissmann,
Institut für Molekularbiologie I, Universität Zürich,
8093 Zürich, Switzerland.

NOV 1 REC'D

Under conditions designed to give selective denaturation of DNA, endonuclease S₁ specifically and efficiently excises a DNA insert from a hybrid at the poly(dA)-poly(dT) junctions. The hybrid plasmid PBG (Maniatis et al., Cell 8, 163), 200 µg/ml, was digested in 45% w/v formamide, 0.2 M NaCl, 1 mM ZnSO₄, 30 mM Na acetate (pH 4.5) with 600 units/ml S₁ endonuclease (Wiegand et al., J. Biol. Chem. 250, 8848) for 30 min at 50°C. This treatment cleaved the DNA into only two fragments. The smaller one, with an average of 575 base pairs, was somewhat heterogeneous and contained all of the globin specific sequences.

Since S₁ endonuclease is rapidly inactivated under the incubation conditions described above, the concentration required for optimal cleavage (about 10 times that which would be required in the absence of formamide) must be determined experimentally. Underdigestion results in a somewhat larger, more heterogeneous product, while overdigestion reduces the size of the globin DNA fragment and causes an additional cleavage in the larger DNA piece.

Several viral DNAs, in particular λ, φ29 and T7 DNA gave defined and characteristic cleavage patterns upon S₁ endonuclease digestion at optimized formamide concentrations.

(Bioch. Biophys. Acta, 1976, in press)

December 1976
Memo NAR-65/1

Generation of new restriction sites by mutation in bacteriophage T5

John Davison and Françoise Brunel

OCT 16 REC'D

MRC Unit of Virology, University of Glasgow.

Present Address: Beatson Institute for Cancer Research, Garscube Estate, Glasgow, G61.

Bacteriophage T5 is insensitive to in vivo restriction by the Eco RI restriction system, although the DNA has six Eco RI sites in vitro. We have isolated mutants of T5 which are sensitive to in vivo Eco RI restriction and have genetically mapped these mutations in the First Step Transfer (FST) portion of the genome (i.e. that portion of DNA which enters the cell in the first 10 minutes after infection).

Tues. Feb. 15th

Grants all done! Meeting attended!

Probe for direct Hb clone detection





If cloned Hb is $\sim 1\%$ of λ , then it would need $100 \times$ cold λ to $\downarrow$ background to equality - assuming no cross-linking. Better to pre-purify cloned piece prior to labeling. Since there are no R1 sites left & Hb2 leaves still $\sim 5\%$ λ DNA - need to try S1 nuclease on $\Delta A/\Delta T$ tails.

$\therefore$ try S1 / formamide on Hb clone e.g. ~~4~~ 4
or "Hb" 3 clone ($\Delta A/\Delta T$)
or 3A Δ lac

Add "stuffer DNA" $\sim 1 \mu\text{g} / 20 \mu\text{l}$ R⁺ mix
(Harvey has 10-113 W Ch 3A Δ lac at $0.66 \mu\text{g}/\mu\text{l}$ in DNA dialysis for 3A Δ lac. For stuffer use "Ch 3A/R1 10-23" at $\frac{1}{2} \mu\text{g}/\mu\text{l}$ in d.w. Heat before use 5 min at 100°C)

10x S1 buffer (Harvey 9-131) - ~~filter~~ sterilized
2M NaCl
0.5M NaAc
10 mM ZnCl_2
pH 4.5 with NaOH

Dilute S1 in 50% glycerol. Available is: HF 10-19 ex JR 1 μl of
(Harvey has some) which works on $10 \mu\text{g}$ DNA in 1 ml R⁺



Marker is cB2 KH100 min 5

61

3A Hb4 OD = 3 i.e. $\sim 0.15 \mu\text{g}/\mu\text{l}$ (label $\phi 61$)

3A Δ Lac 10-113 W at $0.66 \mu\text{g}/\mu\text{l}$

3A Hb3 at $0.65 \mu\text{g}/\mu\text{l}$ (m93)

Test: -

	($\frac{1}{10}$)	($\frac{1}{3}$)	(1)	(3)	(Hb4)	(Hb3)
Deionized formamide	9 μl	9 μl	9 μl	9 μl	9 μl	9 μl
10X S1 buffer	2 μl	2 μl	2 μl	2 μl	2 μl	2 μl
Stuffer DNA (Hb4)	2 μl	2 μl	2 μl	2 μl	2 μl	2 μl
D.W	3 μl	3 μl	3 μl	1 μl	-	3 μl
Ch 3A Δ Lac	3	3	3	3		
3A Hb4					6	
3A Hb3						3
S1 nuclease	1 $\frac{1}{10}$	1 $\frac{1}{3}$	1	3	1	1

13 μl of
63 μl +
14 μl 10X
+14 μl HF9-131

2.22 pm to 50°C (air) 3.0 pm to ice.

n21 Hb3 BDM

New Marker ~~n21 Hb3 BDM~~ $\phi 60$ (by mistake in 20X Hening)

3x S1 3A Δ Lac

Hb3 1x S1

Hb4 1x S1

1x S1

$\frac{1}{3}$ x S1

$\frac{1}{10}$ x S1

3A Δ Lac

1.0 μl $\sqrt{\text{cm}}$

4x Hening

0.7% Agarose

n21 Hb3 BDM $\leftarrow \phi 61$

off 10 pm

A possible faint band on Hb4. test with probe and repeat without "stuffer" DNA


$\rightarrow$ (over)

10.40 pm to sl^m I (1.5m WAG 0.5m NaOH)

10.55 pm

-- sl^m II



11.10 pm ~~to transfer~~ (3m WAG 0.5m Tris)



11.20 pm away

80° after 2xSSC rinse

Next am. dried v2 for at



Friday Feb 18th

Repeat S1 tests without "stuffer" DNA

	$\frac{1}{10}$	$\frac{1}{3}$	1	3	Hb4	Hb3	
Deionized Sawmillside 10xS1	Hpl	Hpl	Hpl	Hpl	Hpl	Hpl	Deionized Sawmillside 63pl [+14pl] 10xS1 buffer ✓
P.W.	5✓	5✓	5✓	3✓	2✓	5✓	
Ala3A Alac 10-113W	3✓	3✓	3✓	3✓			
3A Hb4					6✓		
3A Hb3						3✓	
S1 nuclease	1 of $\frac{1}{10}$	1 of $\frac{1}{3}$	1	3✓	1✓	1✓	

2. 25pm to 50° ± 2° (trouble with thermostat)

$\frac{1}{2}$ hr later to ice

On 4.45 pm 1.0 v/cm 4X Heming 0.7% Agar

Marker mix
Alac 3xS1
Hb3 1xS1
Hb4 1xS1
Alac 1xS1
Alac $\frac{1}{3}$ xS1
Alac $\frac{1}{10}$ xS1

off 11.45 pm

Conclusion: Same piece in Hb4
It is ~ 3.4% !

17
14.2237
14.22Saturday Feb 19thPreparation of probe

If the $\phi 61$ piece is correct can use it to tell Hb^+ from Hb^- simply. Rather than prepare it $\rightarrow$ label (which saves label but is more wasteful of DNA + time), I will label it + isolate. This will give enough for test.

Suppose it is really 1% of λ , & that at $15^\circ C$ I can get 10% incorpⁿ. Then 0.1% of total label will be in $\phi 61$. With $50 \mu Ci$ this will be $.05 \mu Ci$ or $100,000$ cpm. — enough for a test but it is clearly better to go the other way round finally (i.e. purify $\rightarrow$ then label)

Take 30 μl of dATP | (2nd Feb lot. $1 \mu Ci/\mu l$)
 30 μl of dCTP | 1977
 30 μl of dTTP | i.e. $42.3 \mu Ci$ total
 + 52 μCi total
 60 μl of dGTP | (11th Jan 1977 $1 \mu Ci/\mu l$)
 i.e. $9.6 \mu Ci$

✓
 { 100 μl HF 9-13-3 10x nick translation buffer
 + 100 μl l.w. + 0.7 μl mt
 Note
 → 80 μl to load down (^{32}P) dNTPs ✓
 ✓ 20 μl (3 μg) 3A Hb4 DNA ✓
 ✓ 2 μl DNA polymerase m151
 To $15^\circ C \pm 2^\circ C$ at 2.10 pm (except for brief $20^\circ C$)
 10 μl of 1% DEPC at 3.10 pm

To DNA buffer .01M Tris .001MEDTA pH 8.3
6100 column.

1	<0.1mR/hr	6	38
2	<0.1	7	6
3	<0.1	8	0.9
4	0.45	9	0.4
5	38	10	~ same

Pooled 0.72 ml.

5 μ l in Bray's

gave

on 1.45% gain 74300 cpm
50-1000
E F

$$\text{Total} \therefore \frac{74300 \times 2 \times 72}{2.2 \times 10^6} = 4.9 \mu\text{Ci (close)}$$


enough to 10% with unpoised #7).

Added 75 μ l 3M NaAc + 2.4 ml EtOH
- cooled in dry ice & left at $\sim 20^\circ\text{C}$ from
4:10 pm to 4:35 pm. Cent! Savell (same losses)
Washed 1x with 80% EtOH. Blew down. - difficult to tell - up to 20%


✓ Take up in 60 μ l of { 45 μ l formamide
10 μ l 10x S1 buffer
45 μ l d.w. }
Then to 50°C & + 3 μ l S1 AF 10-19 ExJR at 5:25 pm

5:55 pm Froge.

669 Post
S1



Fast
bulk
+1
2-1
1988
Stoo
bulk



UNIVERSITY OF
THE NORTH
CAROLINA

JAYSON
NOTION

Sun. Feb. 20th

Preparative run

On 12 noon Off 4.15 pm


Deleted probe 069 Post S1	
marker mix	0.7% Agarose
Probe 069 Post S1	Hx4
	2v/cm 67mA
3 little deleted	[Air cooled by fan]
Marker mix	

On to Saran Wrap autorad at 4.25 pm
developed at 4.55 pm for quick look

Pattern dark enough but bands not show,
i.e. stained. Cut out

$\sim 10 \text{ mR/hr}$ → PROBE band (i.e. marker) $\sim 8 \text{ mR/hr}$ $\sim 6 \text{ mR/hr}$
 PROBE +1 and -1 (one with ahead & behind)
 BULK Slow → Fast (to cover autorad. areas)
 Freeze ↑
 $\sim 5 \text{ mR/hr}$ $\sim 16 \text{ mR/hr}$

DNA



Tris 0.06M Mg 0.06M pH 7.2

units / pl

Monday Feb 21st

Probe tests


Can test the five potential probes: —

Hha digest $\alpha 3A$, $\alpha 3A\Delta lac$, $\alpha 3AHb3$, $\alpha 3AHb4$
— transfer — hybridize

	3A	3A Δlac	3AHb3	3AHb4	← (~ 2pg total)	
10x Hha Buffer	7 μ l	7 μ l	7 μ l	7 μ l		
ADA19 (~0.8pg) /pe	25 μ l	—	—	—		
10-13 W (~0.3pg) /pe	—	3 μ l	—	—		
m93 Hb3 (~0.5pg) /pe	—	—	3 μ l	—		
obl Hb4 (~0.15pg) /pe	—	—	—	13 μ l		
D.W.	58 μ l	58 μ l	58 μ l	58 μ l		
Hha enzyme	2 μ l	2 μ l	2 μ l	2 μ l		

To 37°C at 11.07 am → 1.07 pm
Ice

Gel is 95 x 147 mm



2x 95 x 285

2x 95 x 147

milli 110 x 150

On 1.30 pm 0.75 V/cm 95 mA

LTH 1.5% Agarose

Off 11.30 pm Coated 15 min Sol^xI 20 min Sol^xII 5 min

Bands, KI

Sulfate
Polysaccharides

Trm DEAE

6M KI

10mM NaPO₄ pH 7

Load on ^{thin} commercial HA BioRad

Wash with low PO₄ to get rid

KI to get rid of
Bran

of
KI

Lo PO₄ → KI

Batch

100% Infectants on SV40

Hin II on Hin III

Tues. Feb 22nd
 (Transferred as usual)

2x SSC rinse

Bake ~2 h 80°

Overnight 9:30pm → in sterile
 Denhardt's 6x SSC at 70°C with slake.

Wed. Feb. 23rd

Probe extraction

Plan Will try boiling water bath to melt Hemings 4x - containing gel (it will be brown a bit from staining). + equal vol 4xH. Then add 1M NaOH to give 0.3M followed by 25 2M Phos pH 7 neutralizing buffer (equal vol.). Then add 6xSSC.

(Test
P-1)

Took out frozen probe - 1. highly centrifuged after thawing. Recovered

0.6mR/hr
1.4mR/hr

581 mg liquid } 876 mg total.
295 mg solid }

↑

Added 0.8 ml ~~0.8 ml~~ 4x Hemings (pH 8)
Boiling water < 5 min. - dissolved.

Added back liquid to ~1.5 ml total

Cooled ~~Added 0.3 ml 2M NaOH~~ & it re-set

& or re-melted & added 1.5 ml more 6xSSC

Added 0.5 ml 2M NaOH pH 12 ~~pH 12~~ ~~Annealing~~

∴ 0.3 ml more

Transferred to 1.6 ml neutralizing buffer
pH close to 7 & semi-solid. Use it all.

✓

Test P+I

0.79 ~~g~~ total

Added 2.5 ml 6xSSC (Benhardt) Annealing
Buffer (pH 8-9 - closer to 9)

Added 0.5 ml 2M NaOH pH > 13

Added 1 ml neut.⁹ buffer (use a little less)

-- back titrated to pH 7.5.

Set up.

Probe test

0.72

Added 2 ml 6xSSC Annealing
Buffer

Added 0.5 ml 2M NaOH

Added 0.8 ml neut.⁹ buffer - pH 7

-- added 0.1 ml 2M NaOH - pH 7.5


- - - 0.1 ml - pH 9

- - - 0.025 ml neut.⁹ pH 8.5



Used $\frac{1}{2}$ for test - Troze test

as 0.79 P₀ PROBE

To 70°C at 1.15 pm Wed → ~9:30am Thurs




With cDNA probe
at 70°C only one
Hha piece can be found
See also, 0/30



Thurs. Feb 24th


Probe test (contd.)



P₀ off.

Used
P₀ #81

probe saved -
it is yellowish.



P-1



P+1

0.5% SDS
Rinsed all in ~100 ml 2xSSC →
to 70°C (together) at 10.45am

→ 11.45am 11.55am → 12.25 1230 → 1.45pm

Rinsed in 2xSSC. Air dried → to autoraf.
at 2.20 pm Off 8.30am 9.0am → 9am
Th. Fri Fri Mon

Conclusion: Agarose method v. unimpressive but
probe is fantastically specific. ∴ repeat with SI
first. Try old DEAE-HC method to clean up agar.

Screening λ gt Recombinant Clones by in situ

Hybridization to Single Plaques

David Benton^{*} and Ronald W. Davis

Department of Biochemistry

Stanford University School of Medicine

Stanford, California 94305

* Dept. of Genetics and Cell Biology
250 Biological Sciences Center
1445 Goodwin Ave.
St Paul, Minn 55108

limited only by the availability of complementary nucleic acid
which can express foreign genetic information. The third method is
limited to genes for which appropriate mutations are available in hosts
complete nucleic acid sequence of any gene. Application of the second is
first method is limited by the practical difficulty of physically purifying the
DNA probes to screen clones produced through a shotgun experiment. The
genome of the donor organism (3); or (iii) using complementary RNA or
(ii) genetic selection of the desired gene after shotgun (2) cloning the
the gene or its transcript before insertion into a vector for cloning (1);
isolation has been accomplished in any of three ways: (i) purification of
isolate large quantities of DNA corresponding to any single gene. This
Recombinant DNA technology has made it possible, in principle, to

ABSTRACT

A rapid, direct method for screening single plaques of
 λ gt recombinant phage is described. The method allows at least
 10^6 clones to be screened per day and simplifies physical containment
of recombinants.

Recombinant DNA technology has made it possible, in principle, to isolate large quantities of DNA corresponding to any single gene. This isolation has been accomplished in any of three ways: (i) purification of the gene or its transcript before insertion into a vector for cloning (1); (ii) genetic selection of the desired gene after shotgun (2) cloning the genome of the donor organism (3); or (iii) using complementary RNA or DNA probes to screen clones produced through a shotgun experiment. The first method is limited by the practical difficulty of physically purifying the complete nucleic acid sequence of any gene. Application of the second is limited to genes for which appropriate mutations are available in hosts which can express foreign genetic information. The third method is limited only by the availability of complementary nucleic acid probes and has been the most widely applied.

Screening methods have been developed for both recombinant plasmid (4) and bacteriophage (5, 6) cloning systems. The most rapid of these, the methods of Grunstein and Kramer, are based on a technology developed by Olivera and Bonhoeffer (7) for growing and lysing cells on membrane filters. The speed and simplicity of these methods is limited by the need to pick and spot plaques or colonies from plates onto filters for further growth. We have found that a single plaque of a λ gt recombinant phage contains enough phage DNA for detectable hybridization to complementary labelled nucleic acid and that this DNA can be fixed to a nitrocellulose filter by making direct contact between the plaque and the filter. These findings have allowed us to develop a method for screening

recombinant phage which has significant advantages over previous methods:

(i) Physical containment of recombinant molecules (and organisms carrying them) is made simpler because this method requires fewer manipulations of viable phage, requires the growth and handling of much smaller quantities of recombinant DNA-containing phage and cells, and produces no contaminated materials other than the petri dishes on which the plaques are grown. (ii) At least 2×10^4 plaques can be screened per hour (8), which compares with about 100 per hour by the fastest previous method (6). (iii) Because up to 2×10^4 plaques can be screened on a single petri dish and nitrocellulose filter, the method requires smaller amounts of materials, including radioactively-labelled probe. (iv) Because screening can be done directly from the transfection plate, all the independently-constructed phage containing similar or identical inserts can be isolated and individually characterized. This can give information on the heterogeneity of repeated genes or of genes from different somatic tissues.

We believe that this method will be widely applicable and we, therefore, present it here in detail. Preliminary experiments and the model system used for optimization are also described.

Cells (9) are infected with phage, plated in L-soft agar on L-plates, and grown for at least 12 hours at 37°C . Large plaques are usually desirable and 0.1 ml of late log phase cells per plate has been found to give large plaques of many λ gt phages. The number of phage which can be plated on a single petri dish depends on the relative growth rates of the desired recombinant and the background phages (see below). After growth, the plates are placed at 4°C for about 15 min to harden the agar. Phage and DNA are transferred to a nitrocellulose filter by placing the dry filter (88 mm, Millipore HA or Schleicher and Scheull, BA 85) on the lawn, being careful to form no air bubbles between the soft agar and the filter. Phage are allowed to adsorb to the filters for 1-20 min. If duplicate

recombinant phage which has significant advantages over previous methods: recombinant phage requires fewer manipulations (and organisms carrying recombinant phage)

filters are desired, several filters can be adsorbed sequentially for 30-60 sec each or filters can be stacked and adsorbed at once. If filters are stacked, they must be allowed to adsorb for at least 20 min. During adsorption, it is convenient to mark the filters and plates for orientation (see below). Following adsorption, the filter is carefully lifted from the plate to avoid removing the soft agar layer which occasionally adheres to the filter. Should this occur, the agar can be removed by gently shaking the filter during the denaturation and neutralization steps. DNA and phage are denatured and fixed in situ by dipping the filters in 0.1 N, NaOH, 1.5 M NaCl for 20 sec and the filters neutralized by dipping in 0.2 M tris, pH 7.5, 2 x SSCP (11) for 20 sec. Filters are blotted and baked at 80°C in vacuo for 1.5 to 2 hours.

Hybridization to the nitrocellulose replicas is carried out in a siliconized glass petri dish using $10^5 - 10^6$ cpm of ^{32}P -labelled probe (10) per filter in enough 5 x SSCP, 50% formamide to cover all filters. Hybridizations are typically for 12-18 hours. Buffer and probe can be recovered and reused for at least two weeks. Following hybridization, filters are washed in a large (10-15 ml/Filter) volume of 5 x SSCP, 50% formamide at 42°C for 30 min. and then for 20-30 min in 2 x SSCP at room temperature. The filters are then covered with plastic wrapping film and placed against X-ray film for autoradiography. 24-48 hours of exposure is usually sufficient with a homogeneous probe of specific activity of $10^5 - 10^6$ cpm/mg. X-ray intensifying screens (Kodak X-Omatic

Regular) have been used with flash-activated film at -70°C to shorten exposure time.

The method described above should be generally applicable to all recombinants of λgt phages. The experiments described below used the phage $\lambda\text{gt-Sc 1109}$ (6, 12) in model systems to optimize the method. Phage grown on E. coli ~~DX~~ C600 for less than 9 hours give negligible autoradiographic signals; 12-14 hours of growth is optimal. Although T-plates generally yield larger plaques, these apparently contain fewer phage particles and less DNA than comparably sized plaques grown on L-plates, since, at all growth times, nitrocellulose replicas of T-plates gave lower signal than those of L-plates (13).

To determine the maximum number of plaques per plate which can be screened under various relative growth conditions, two experiments were performed. In the first, to model a situation in which the background grows much better than the desired recombinant phage, a small number of $\lambda\text{gt-Sc1109}$ phage were grown together with increasing numbers of λgt2 (14) phage. No effect on signal was apparent up to about 2000 pfu/plate, but the signal was weak and variable with more than 4000 pfu/plate. In the second competition experiment, $\lambda\text{gt-Sc1109}$ was plated with increasing numbers of $\lambda\text{gt}^{\text{+}}\text{-}\lambda\text{B}$, which has approximately the same plaque size as the recombinant phage. In this case, sufficient signal was obtained with more than 2×10^4 pfu/plate. A practical rule seems to be that the method will accommodate the greatest number of plaques that does not produce confluent lysis.

Time of phage and DNA adsorption to the nitrocellulose replica filter does not seem to be critical; adsorption times of 1-60 min have

been used and little difference noticed. Very long adsorptions resulted in slightly more diffuse plaque-spots on the nitrocellulose replicas and the autoradiograms. Stacks of at least five filters can be adsorbed at one time to produce replicates (for hybridization to several different probes, for example). For most applications, better replicates are produced by sequentially placing the filters on the plate for a brief adsorption to each (Fig. 1).

Dipping the nitrocellulose replica directly into a beaker of denaturing or neutralizing solution does not result in loss of signal or in distortion of the shapes or relative positions of the plaque spots. Denaturation and neutralization times of 0 to 60 sec all give acceptable results; 15-20 sec of each is optimal, giving an approximately 2-fold increase in signal over either no denaturation or no neutralization. Baking the filters before hybridization is not necessary, but in this model system the signal was increased approximately 2-fold by baking for 90-120 min.

Following autoradiography, it is necessary to align the autoradiogram of the nitrocellulose replica with the original plaque plate to allow picking of the plaques corresponding to positive signals. The most reliable system we have used for this alignment involves labeling the filters and plates during adsorption by puncturing the filter at 3 points around its circumference with a hypodermic needle and injecting 0.1-0.2 μ l India

1. M. Thomas, R. L. White, and R. W. Davis, Proc. Nat. Acad. Sci.
USA 73, 2294 (1976);
B. E. Noyes and G. R. Stark, Cell 5, 301 (1975);
S. G. Clarkson, H. O. Smith, W. Schaefer,
Birnstiel, and M. A. S. Schaefer.

REFERENCES

ink into the agar at each point. The ink diffuses very little during the 2 days normally required for hybridization and autoradiography and allows absolute orientation of the filters, autoradiograms, and plates. From a plate with more than 1000 plaques, we normally pick about 20 plaques from the area of each positive signal for rescreening and plaque purification. Once pure clones are obtained, the sizes and restriction spectra of the inserted DNAs can be rapidly determined by the method of Cameron, et al (15).

The large capacity and speed of this method make it practical, we believe, to isolate a eukaryotic unique gene by screening all the recombinants produced through shotgun cloning the entire genome. This should make possible isolation of a large number of genes for which physical or genetic purification schemes are not available.

2 days normally required for hybridization and autoradiography and
 the ink diffuses very little during the point. The ink diffuses very little during the
 and plates.

REFERENCES

1. M. Thomas, R.L. White, and R.W. Davis, Proc. Nat. Acad. Sci. USA 73, 2294 (1976);
 B.E. Noyes and G.R. Stark, Cell 5, 301 (1975);
 S. G. Clarkson, H.O. Smith, W. Schaffer, K. W. Gross and M.L. Birnstiel, Nuc. Acids Res. 3, 2617 (1976).
 G. N. Yenikolopov, A.P. Ryskov, T. Nitta, and G.P. Georgiev, Nuc. Acids Res. 3, 2645 (1976);
 S.C. Hardies and R.D. Wells, Proc. Nat. Acad. Sci. USA 73, 3117 (1976);
 R. Higuchi, G.V. Paddock, R. Wall, and W. Salser, Proc. Nat. Acad. Sci. USA 73, 3146 (1976).
2. A shotgun experiment is defined as one which involves the production of recombinant DNAs between the vector and total or partially purified DNA from a cellular source.
3. K. Struhl, J.R. Cameron, and R.W. Davis, Proc. Nat. Acad. Sci. USA 73, 1471 (1976);
 L. Clarke and J. Carbon, Cell 9, 91 (1976).
4. L. Kedes, A. Chang, D. Houseman, and S. Cohen, Nature 255, (1975);
 M. Grunstein and D. Hogness, Proc. Nat. Acad. Sci. USA 72, 3961 (1975).
5. K. Jones and K. Murray, J.M.B. 96, 455 (1975); Yu. V. Ilyin, N.A. Tchurikov, and G.P. Georgiev, Nuc. Acids Res. 3, 2115 (1976);
 B. Sanzey, O. Mercereau, T. Temynck, and P. Kourilsky, Proc. Nat. Sci. USA 73, 3394 (1976).
6. R.A. Kramer, J.R. Cameron, and R.W. Davis, Cell 8, 227 (1976).
7. B.M. Olivera and F. Bonhoeffer, Nature 250, 513 (1974).
8. This calculation is based on the worst case in which only 2000 plaques can be grown on each of 50 plates, the screening of which would require about 5 working hours over a three-day period.
9. *E. coli* C600 $rk^- mk^+$ was the usual host. Cells and phage are described in J. R. Cameron, S.M. Panaseuko, I.R. Lehman, and R.W. Davis, Proc. Nat. Acad. Sci. USA 72, 3416 (1975).
10. In the model experiments described, the probe was total yeast rRNA (^{32}P -labelled) prepared by the method of G.M. Rubin, Methods in Cell Biol. 12, 45 (1972).

11. SSCP is Standard Saline Citrate Phosphate, 1 x = 120 mM NaCl, 15 mM NaCitrate, 13 mM KH_2PO_4 , 1 mM EDTA, titrated to pH 7.2 with NaOH.
12. λ gt-Sc1109, the gift of R. Kramer, is a recombinant λ gt phage which carries 2 EcoRI fragments of *Saccharomyces cerevisiae* DNA, one of which carries the 5.8 S and 26 S rRNA genes. Approximately 5% of this phage's DNA is complementary to rRNA.
13. T-plates = 1% Bacto tryptone, 0.5% NaCl, 1% Agar, made to pH 7.5 with NaOH.
L-plates = 1% Bacto tryptone, 0.5% Bacto yeast extract, 0.5% NaCl, 1.25% Agar, made to pH 7.5 with NaOH.
soft agar in each case is 0.7% in Agar.
14. λ gt2 [λ gti in Struhl, et al (3)] and λ gtAB were the gifts of M. Thomas.
15. J.R. Cameron, P. Philippsen, and R.W. Davis, submitted to Nuc. Acids Res.
16. We thank T. St. John for helpful discussions and Lynn Horn for help in manuscript preparation.
17. This work was supported by NSF Grant PCM 76-02600 and NIH Grant GM-021819-03.
One of us (WDB) has been supported by NIH Training Grant 5T01-GM 01156 and by a fellowship from the Bush Foundation, St. Paul, Mn.

Fr Feb 26 ch

Clean up of agarose



DEAE 23 A branch fibrous ex.
Whatman Acidified with $\sim 1M HCl$
 $\sim 2M NaCl$ & washed with d.w.

Recycled with $0.5M NaOH$ & d.w.
& repeated once more. Finally
acidified ^{to pH 1} with HCl alone, d.w.
washed } & dried at 42° until
to suspended } Monday am,
pH ~ 4

0.7% agarose made as usual with 4x Hemmings
0.75% agarose + 50 ml d.w. (+ 1.5g moist
DEAE-HCl after first load) boiled. Milled
0.22. Added 20x buffer & d.w. to usual.

RI digest of 3A available from 13 (has
 $\frac{14}{23} \mu g/\mu l$ in 2.6 $\mu g/\mu l$) — not enough!


~~0.22 + 40 μl HX buffer~~



New Digest: —


{ 20 μ l ADA 19 ch3 (+lac) DNA ($\approx 14 \mu$ g DNA)
+ 2.3 μ l 10X R1 + $\approx 0.4 \mu$ l DGR1
to 37° at 1.40 pm $\rightarrow$ 2.30 pm

Diluted with 50 μ l L.W.



First cut off origin 2 cm. from each
o stained. Then edges. Then correct
band. Probably OK - but tended to over-
estimate migration.

1 ~~0.2.45 pm~~ 5ml two pieces washed in 1X R1 buffer
with gentle rock for 15 mins to get to
R1 salts (? in real test run get in ^{nick translation} salts)
 $\approx 1.15 pm$



$\phi 1$
 2pl / 1pl 10XR1
 L Lot 2 8pl ~~10pl~~
 d.w

Him
 Super
 Sam

$\uparrow$ to $6\frac{1}{2}V$
 5.10 $\downarrow$
 5.30
 On 4.50pm

D-E	3v/cm
D+E	Hx4
U+E	0.7 ⁹ /atgavok
U-E	

Decanted liquid - Freeze

Mon Feb 28 ch

Contⁿ of DEAE testThawed gels & collected ^{~200} ~~20~~ μ l of super^t from ~~2 test tubes~~ "usual" ($\sim 10 \mu$ l was lost) $\sim 300 \mu$ l from "DEAE"

Refreeze gels

Refreeze super^s

Tested on supernatants: -

	DEAE	DEAE	USUAL	USUAL	
Hind III	-	+ 2 μ l	-	+ 2 μ l	
Super ^t Sample	40 μ l	40 μ l	40 μ l	40 μ l	11:10 am to 37°
					↓
					11:40 am

 $\sim 20 \mu$ l of each ^{iced} ~~iced~~ - rest run

On 11:50 am

Off 12:00 pm

0.7% Agarose

H x 4

3 V/cm ~ 100 mA

Rerun 1:40 pm

↓
2:40 pm1:30 pm back to 37°
↓
4:30 pmNo digestion
in dry in / hr

DEAE - eng

DEAE + eng


USUAL + eng

USUAL - eng

 $\uparrow$ to 6.2V
 $\downarrow$
to 5.50

Agarose

Odd result !



Mon. Feb 28th

Nick translation from gel

Decided to try: —

Can titrate any gel
troubles and
more enzyme.

Run RI of $\alpha 3A$ to get lac.
Freeze gel after dialyzing to nick trans.
buffer (unless I can run in it).
Add dNTP's (P₂P), DNA pol. incubate.
Add cold ~~to~~ pyrophosphate 6RSSC etc.
Denature + hybridize.

12:20 pm

10 μ l ADA 19 $\alpha 3$ (+lac) DNA ($\approx 7 \mu$ g DNA
 $\approx 0.7 \mu$ g lac)

+ 1 μ l 10x RI ($\phi 1$) + 20.2 μ l BGR1

off 1:10 pm to ice. Diluted with 110 μ l 1x Tris/EDTA

Nick translation buffer is

0.05 M Tris pH 7.8

Make
to take care
of EDTA

5 mM

1 mM ME

50 μ g/ml BSA

Mg $\alpha 2$

0.001 M EDTA
Run in this with EDTA


Add these as 10x later

0.05 M Tris / 0.001 M EDTA



0.7% Agarose / DEAE beads

pH 7.8

on 3 pm 1v/cm off 8:15 am



Offer still



Tues. March 1st

Nick translation again

Contaminant - 0.5M Tris HCl / 0.001M EDTA
 being so poor on e.p. but can
 go back to Henning's 1X :-

• 0.4 M Tris - acetate
 • 0.2 M Na - acetate
 • 0.02 M Na₂ EDTA
 • 0.18 M NaCl

pH 8.05

I adjust it to pH 7.8. When diluted to
 1/2 X with equal vol. of d.w. containing 2 mM ME
 100 µg/ml BSA
 9 mM MgCl₂



it sh^d approximate nick translation buffer

i.e. repeated digest { 10 µl ADA 19 Charon 3A
 2.0 µl 10x RI (pH)
 3.15 µl (R.T.) to 37° ~ 0.2 µl DG RI

Remake 0.7% Agarose / DEAE treated
 in 1X H

0.7% HX1
 DEAE treated
 4 V/cm
 150 V, 20 min

del. 2nd 11 2nd



Wed - March 2nd

Repeats of recent gels

ADA 19 ch 3A	10 pl	} 10.30am to 37°
10x RI buffer	12 pl	
10 RI XXXXXX	NO. 4 pl	
S.W.	100 pl	

1.30pm 2.4v/cm in following systems: -

0.7% Agarose	DEAE-treated	Tris .05M / 0.01M EDTA
-----	DEAE-treated	Hx1
-----	Not treated	Hx1
-----		Hx4

Off 8.0pm

Caution DEAE is trouble if buffer is not high.

∴ best ~~DEAE~~ Tris . HCl / EDTA straight.

— but meanwhile best available Hx1

On 5.25pm

Off 9.0am

Cut out band for lac (330mg) & freeze

Test regions	
{ Sample	Hx1
	0.7% Agarose
	1v/cm

Thurs March 2nd

Further nick translation idea

Since sep. is poor at low salt, it does not look good to try the original idea,

- i.e. gel ep., freeze, nick translate, hybridize

This is ok only if probe is already hot i.e.

gel ep., hybridize. $\therefore$ try Kelly's procedure

(see p 74) which is to dissolve gel in 6M KI

10mM Phos. pH 7 & a bead to small Hydroxyl
Appetite column. Wash with KI to ensure gel
& with low phos. to get rid of KI.


Then elute with high phos. Salt would need
to dialyse or Sephadex

Before going up my own idea, try adding hot
KI to (frozen) gel & running over small Sephadex
column with translation buffer in it. As expected
- it clogs.

$\therefore$ test: — (1) vol (3) vol

Add to e.g. 300mg gel (3x300 μ l of 6M KI
10mM Phos. pH 7. Add 30mg HAP (0.1 vol)

(Capacity is 1mg DNA / 1ml swollen gel) Wash 1x (by centⁿ) with KI
 $\therefore \approx 30$ pg



So it works
Need 32 for yield

~ 50 μ l

Wash 2x with 10mM phos. Take up in 0.6M phos. to give
 0.3M $\rightarrow$ apply to top of small Sephadex against ~~red blood~~
 1mM EDTA buffer. Collect using precalibrated volumes. Ppt.
 with Ethanol if reqd.

Q95 has gel (330mg) + 1ml $\sim$ 6M KI KPhos
 $\sim$ 10mM. (cat?)
 + add DTT to
 keep I_2 down

+ 33mg try HTP (BioRad)

$\uparrow \uparrow \uparrow \uparrow$
 (Slide at 1st)

Vortexed well. left $\sim$ 10min

— — — again. Cent'd. Decanted.

Washed 1x with KI/phos. Cent'd. Decanted.

Washed with 10mM KPhos 2x

Eluted with 650 μ l 1M $\left. \begin{matrix} K_2H \\ KH_2 \end{matrix} \right\}$ Phos.
 (Final vol. $\sim$ 150 μ l)

On 4:15pm $\rightarrow$ 5:15pm

0.7% Agarose

4xH

3V/cm 100mA (max)

Repeats

On 5:20pm

0.7% Hx4

1V/cm

9:20am

ADA 19 Ch3A 10 μ l

10xRC 12 μ l

DGR1 $\sim$ 0.4 μ l

D.W. 100 μ l

4:25pm to 37°

$\downarrow$
 5:10pm

Friday March 4th

1.01g sample from lac gel 099
frozen & thawed

+2 ml Frisch 6M KI 10mM $\text{KH}_2/\text{K}_2\text{H Phos.}$ / 0.1mM EDTA
(sat. is slightly less)

$\approx 6500 \text{ cpm}$

Added to it (for tracer purposes) 2 μl of
~~Ch 3A~~ LAF nick-translated Hha
Ch 3A Δ lac which is $\approx 1.5 \times 10^6 \text{ cpm}/\mu\text{g}$
- so only $\approx .004 \mu\text{g}$ total ~~Ch 3A~~

12 noon $\approx$ 30mg HTP obtained from 30mg washed 2x
with 1g settle with KI / Phos.

left rocking until 2pm
Washed 2x KI

2x 10mM Phos to vol. 150 μl

Added 150 μl 1M $\text{KH}_2/\text{K}_2\text{H Phos}$

Sephadexed to 0.001M EDTA

Salt came in tubes

$\therefore$ pooled 2 thru' 7 (can't find ^{32}P)

Missed! Salt ppt'd $\therefore$ repeat
with more ^{32}P

$108 \equiv 324$
 $72 \equiv 216$
 $36 \equiv 108$

Sat. March 5th

Recombination


It is clear that the SI method on AT is not v. good.

It is also clear that ~~double label~~ nick transⁿ on an SI piece is not going to be easy.

∴ alternate generations.

Take out putative β chain ~~clones~~ clones.

Cut with Hae III :-




Then reclone in EK1 plasmid with no homology to lambda using R1 site cleaned up with SI & but ligated. Detect reclone with cDNA from mouse Hb

Reclone in plasmid & label
with cDNA from globin

If either Hae III site is lost it
will still work on piece 2 or 3.

Present maps are: —

Hb 11



Wed. March 9th

Nick translation of Hb4

Direct n.t. of whole λ , followed by Hha, Bam, Hae etc. c'd be used for piece. Hershey will do it via Hha piece nick translation.

Better current method is to use hot λ DNA Hb via Southern onto cold Hb4 digested with Bam, Hae, & Bam+Hae if m RNA is available.

For either procedure dry down 2 NTPs.
(Find that m RNA not available for 10 days)
 $\therefore$ proceed to n.t.
^{2 1/2 half hrs}
17 1/2% level (Feb 2nd) 2 ATP, dCTP, 2 TTP
available (as excess) 2 GTP not available

Make 75 μ C, if poss.

$\therefore \frac{100}{17.5} \times 25 \mu$ l of each w'd be needed i.e. 142 μ l

142 μ l of each dCTP, 2 TTP, 2 ATP dried

~~142 μ l of each~~

Each is 200 C/mMol

$\therefore 142 \mu$ l (back label $\approx 142 \mu$ C) $\approx \frac{71}{142} \times 10^4 \mu$ M

1 μ g of λ DNA is $\frac{1 \times 10^6}{330} \mu$ M of each base $\approx 710 \mu$ M
or 2 μ g really needed.

take up brief from dNTPs in

90 μ l $\leftarrow$ HF 9-133 ^{1x} nt. buffer $\checkmark$
 (80 μ l stock 10x + 800 μ l w + 0.7 μ l md) $\checkmark$
 + 10 μ l 3AHB4 DNA (1.5 μ g)
 + 0.1 μ l 20mM dGTP (2000 pMoles)
 + 2 μ l DNA pl. m151

$\rightarrow$ To 15°C

1.20 pm

1.20 pm + 10 μ l 1% DEPC

Sephadexed to 0.001M EDTA pH 7.8

	mR/hr		mR/hr		mR/hr
1	<0.1	4	13	7	1.7
2	<0.1	5	17	8	1.1
3	<0.1	6	4	9	1.1



0.10 μ l
³²P ChHb4

Pooled Vol. $\sim$ 590 μ l 5 μ l $\equiv$ 32,000 cpm

$\therefore$ incorp. $\sim$ 1.6 μ C

1 μ l $\equiv$ 6,000 cpm

hesley has Ch3A at 18,000 cpm/ μ l



HLA
2lac
Hb 4

Thurs. March 10th

Hha Digests

			20pl	20pl	20pl	Rest
10x Hha buffer	✓	8 pl				
ADA 19 (3A)	✓	2.5 pl				
10-113W (Δlac)	✓	3 pl				
0109 ^{32P} Hb4		—		6pl	12pl	
hexby ^{32P} (Δlac)		—	2pl			
D.d.w.	✓	53 pl				
Hha en. Batch 1390 labels	✓	4pl				
To 37°		1.30pm	→ 2.40pm			

On 2.50pm Off 9.30pm

D.W.
 Cold mix (residue)
 Hb4
 3A Δlac
 2x Hb4
 Residue ^{32P} labels
 D.W.

1.5% Agarose
 Hx4
 1.5 V/cm 60mA

Stained
 Foxy & exposed
 10pm → 2mon
 Th 24hr Fri
 Fri — Sun
 Noon 48hr Noon

Product is adequate /

Mon. March 14th

Re-alternating vectors

No reason not to do this with
histones too. We have pSC101 histone
clones — probe? ∴ go to λ immediately
for R₁ clones from genome. Can test
the human & mouse R₁ clones also
in the same way. ∴ go to 4A to
get selection v. screening — except length of
repeat is prob. too small for 4A. ??
∴ maybe we should use a non-cross
reacting plasmid with no homology. With
suitable DNA ~~conc.~~ conc. the self-closing
can however be kept low. We need
to calculate efficiencies & total DNA
amounts & in vitro packagingments
& pre-screen group

After discussion: Every reason to pick pBR322, — it is
EK2 in X1776 & can be used semi-selectively
to but-end ligase. We probably will find it best
to stay with λ for all genomic clones, & use
plasmid only for probe preparation once cut-out-able
pieces are in use. It is also probable that
spheres + DNA → 1 round lysis (100 copies) will
be best, since we can plate up to 20,000 phage
but only 1000 sphere plaques/plate, also v. ~ 2000
bacterial colonies/plate.

Cut out putative land from Hb4. Froze.
Rest re-autographed.

Thurs. March 17d

Remake of Hha piece (^{32}P) for Hae III

Make up 1.5% agarose gel with HX1 buffer so as to be able to get Hae III to work after extraction.

Digestion sh't be an ppt! need translated to get plenty of counts in small vol.

~ 300,000 cpm - take 50 μ l Hb4 0109 + add 150 μ l EtOH
— went a little cloudy

~ 900,000 cpm — — — 50 μ l Δ lac D101 B — — —
(hesley) — both in 10^{-3} M EDTA

✓ 2.30 pm to -70°

10x Roberts 60 mM $MgCl_2$ pH 7.9
60 mM Tris
60 mM ME

✓ 2.45 pm spun in Beckman B (10,000 rpm) for 5 min
✓ Blow off ~~EtOH~~ EtOH

✓ Add 2.5 μ l 40 mM $MgCl_2$ to each (to neutralize EDTA)

✓ 20 μ l to each of { 5 μ l 10x Hha i.e. Roberts buffer }
{ 20 μ l Hha enz. Bst 139 (Bst 139) }
{ 40 μ l D $_2$ O }

By finger
there are
more counts
in Hb4 sample!

3.35 pm to $37^{\circ}C$ Off ~ 4.35 pm

off 5.30 to autored
10.30 pm

3 Hb4 product (same on top of gel!!)
3 Δ lac product

on 4.40 pm
1x H 1.5V/cm
1.5% Agarose 12.15 am

Tubes

1	<0.1	5	0.7 mC/hr
2	<0.1	6	0.7 mC/hr
3	<0.1	7	0.1
4	<0.1	8	0.1

Too much label

Just possible that counter was
shortly after

wrong - it quit

I must have muffed one test!

— no — this is not it.

~~A start day!~~
~~for mistakes~~

Fri. March 18th

Remake of ^{32}P plagues

Result not v. good on this piece. . .
remake nick translated products again
- altho' c't done directly from Hae III & select
by cDNA (globin).

Take 2x80 μC total (20 μCi of each dNTP)
per reaction

- today this is 87 μl of each - by error I
took 174 μl . . . save $\frac{1}{2}$ as { Equimolar
Mix dNTP's }
March 2nd lot
~ 400 μl

OK

~~same error here~~
~~If was < 400 μl~~

Take up in 200 μl of

200 μl
4 μl
DNA pol
m151

HF9-133
100 μl Stock 10x
800 μl d.w.
0.7 μl ME

100 μl
70 μl SAHb 4 (~1 μg)

To whole
To next, added

10-113Y (~1 μg)
Harvey's Q3A Δlac
DNA

Can't find it
dammit!!

just do Δlac

To 15° at 12.45 pm.

DEPC (10 μl of 1%) at 1.45 pm
g100 to 10^{-3}M EDTA pH 8

Fri. March 18th.

Test of Hha Hb 4 piece

0119
Agutab
+3A

Marker Standard Hha:—

10X Hha buffer	6 pl ✓
10-113W (Δ lac)	4 pl ✓
ADA 19 (3A)	2.5 pl ✓
D.J.W.	44 pl ✓
Hha eng Batch 139 Biolabs	4 pl ✓

To 37° at 1.20 pm

↓
2.20 pm

+ DEPE to 0.1%

Thawed out 0/15 piece

1 pl gawc
super

126 cpm
-15 cpm

background

~120 cpm

Added 100 pl d.w. - to ice.

3 pm

~~6.20 pm~~

Refreeze stick!

left overnight

5 pl gawc

~ 55 cpm

over background

No good.

1	<0.1	6	12
2	<0.1	7	33
3	<0.1	8	9
4	<0.1	9	~0.5
5	0.16	10	~0.5

Pooled
1,115 μ l
5 μ l 40,300 cpm
4 pC
8000 cpm/ μ l

Passive
Release

Sat. March 19th

Defeat nicktranslation of Δ lac

March 2nd 65

60 μ l of 4 base mix from 0117 tried
to run (i.e. 25 μ C - sh'd be quite
safe)

Took up in 90 μ l $\rightarrow$ { 100 μ l F&I-33
+ 2 μ l 10-1137 stock 10x nt buffer
+ 2 μ l DNAPol m151 800 μ l d.w.
0.7 μ l ME

To 15°C at 1.05 pm DEPC to 0.1% at 2.05 pm

G100 to .001M EDTA

So I poisoned the reaction
with isotope yesterday

0121 Δ lac
8000 cpm/ μ l

Sat. March 19th

Another scheme

Take Hb4, digest with HaeIII & with HaeIII + Bam, run on 1.5% agarose, transfer to Nitrocellulose, hybridize to cDNA & find desired piece. Problem - resolution of 300bp. pieces will be rather poor. Use marker & try it anyway.

Tues. March 29th

HaeIII digests from hAF: —

(2)

Hb4 - P30-31 8pg in ~50pl with Roberts buffer (not inactivated)

(4)

Ch3A Δ lac 6422 7.5pg in ~50pl with Roberts buffer (not inactivated)



∴ digested:

	Hb4 Hae	Δ lac Hae	Δ lac Hae/Bam	Hb4 Hae/Bam	
(m93) 10X Bam buffer	8 pl	8 pl	8 pl	8 pl	
(mT) BAM H-1	—	—	~2 pl	~2 pl	
d.w.	48 pl	48 pl	48 pl	48 pl	
→ (2)	24 pl	24 pl	24 pl	24 pl	
→ (4)	—	—	—	—	
			40 pl	40 pl	
			to 37°	to 37°	
			4:30 pm	4:30 pm	
					5:30 pm / ice


Save 2 for testing

124

3 probable pieces
N. cat. BAM — common to Δ lac.



Should be only
5 BAM cuts.
I can see 3 of
them



Conclusion looks as if
Hae III cuts close anyway to Bam
makes a small piece too small to stick to or find

On ~~6.0 pm~~ off 10.30 pm

1.5% Agarose
Hx4
0.75 v/cm
long gel
33 mA

Hb4	Hae III	} 0.123
Hb4	Hae III / Bam HI	
Δ lac	Hae III / Bam HI	
Δ lac	Hae III	

Stained Photographed. Resoaked in Hx4.
Back on at 0.75 v/cm at 11.10 pm 25 mA
(so not completely back to Hx4, but okay).

off 9 am

Hae III Δ lac digest
is poor


only
cuts.
3 of

~ 9.15	} Sol ⁿ I	{ 3M NaCl 0.5M NaOH
~ 9.30		
~ 9.45		
	Sol ⁿ II	{ 3M NaCl 0.5M Tris pH 7

10.10 am

Transfer started

2 x SSC $\rightarrow$



Flow seems rather fast; I am suspicious of a by-pass.
2 pm Took off towels & replaced with smaller. Can't
see by-pass — but flow became reasonable.
7.30 pm Rinsed 2 x SSC ~ 20 min. Dried. { Oven for 2 hr. 80°C } 10.10 pm
10.25 pm To 70° in Dharf & Solⁿ (cut & on 0.131)

1 < .1	6 < .1	
2 < .1	7 7 mR/hr	} Pooled 1050 μ l
3 < .1	8 33 mR/hr	
4 < .1	9 8 mR/hr	
5 < .1	10 0.7	5 μ l $\equiv$ 36700 7300

0127 7300 cpm/ μ l
 32P Hb4
 nt

Wed. March 23rd


Nick translation of λ 3A Hb4

72 ~~0~~ μ l of 4 base mix from 0117 stored
down (i.e. $\sim 25^\circ\text{C}$)

Took up in 90 μl ✓ + 7 μl Hb4 DNA (1 μg) O.D. 3.5 ex LAF ⁶³¹ 3/77 + 2 μl DNA pol M151	100 μl HF 91-33 ✓ stock 10x nt buffer 800 μl d.w. ✓ 0.7 μl ME ✓
---	--

To 15°C at 11.10 am 0.1% DEPC at 12.10 pm

To 0.01M EDTA by 6:00 Sephadex.



Wed. March 23rd

Hha of labeled Δ lac & Hb4

	Δ lac	Hb4
10x Hha buffer (i.e. 10x Roberts)	20 μ l ✓	20 μ l ✓
0.121 Δ lac 8000 cpm/ μ l	180 μ l ^(1.4x10⁶ cpm) ✓	—
0.127 Hb4 cpm/ μ l	—	180 μ l ^{1.3x10⁶} ✓
(see 072) Hha on Batch 139 Bidsals	2 μ l	2 μ l ✓

To 37° at 1.45 pm; 0.1% DEPC at 2.50 pm

On 3.05 pm ~~at~~ 7.30 pm

~~0.119 Standard digest~~

0.129 ~~0.119~~ Hb4

0.129 Δ lac

1.5% Agarose

Hx4

1.5 V/cm

Stained. Dried down for autorad.

On 8.55 pm → 11.15 pm

11.20 pm → 8.45 am


An important background deactivating trick
 may come out of this accident - in which Multiplier
 went into developer. As far as I recall, it was
 washed in 2 SSC (probably not d.w.) after that. Background
 is very low where developer was there. Try
 developer itself first. Then try pH 11.9 (pH of
 developer) $\rightarrow$ salt conc. of 39 mmo (conductivity
 of developer). Effect is prob. high pH close to but
 less than dissociation of DNA

Typical KOH developer for high contrast

3M) $\leftarrow$ pK 10

1 M)

20M)



to 14

(0.42M)

(0.07M)

(0.22M)

is also 11M

be close to 13

Try 1M NaCl 0.01M NaOH
 - this is pH 11.8 $\rightarrow$ cond. 41 with 50ml
 mmo

or Try 0.4M Na_2SO_3

~~0.2M K_2CO_3~~

0.2M K_2CO_3

NaOH to pH 11.9

Cond. 21
 is 39

An important background decreasing track
 came out of this accident - in which Mullipen
 went into developer. As far as I recalled, it was
 washed in 2 x SS (probably not d.w.) after this. Background
 is ~ zero where developer was there. Try -
 developer itself first. Then try pH 11.9 (pH of
 developer) → salt conc. of 39 (≡ conductivity
 of developer). Effect is prob^{ly} high pH close to but
 not exactly dissociation of DNA

Typical KOH developer for high contrast
 per litre
 D153

25g Hydroquinone	(0.23M)	← pK _a 10
25g ² Metabisulfite	(0.11 M)	
25g KBr	(0.20M)	
50g KOH	(0.89M)	

total ionic strength is ~ 1M
 but this would have a pH close to 14
 Perhaps better formula is

D82

52.5g Na ₂ SO ₃	(0.42M)
8.8 KBr	(0.07M)
8.8 NaOH	(0.22M)

Total ionic strength is also ~ 1M
 but pH would be close to 13

Try 1M NaCl 0.01M NaOH
 - this is pH 11.8 & cond. 41 with 50ml
 in photo

✓ ✓ or Try

0.4M Na ₂ SO ₃	Cond ⁿ is 39
0.2M K₂CO₃	
NaOH to pH 11.9	

Thurs. March 24th

cDNA Hb probe test & gels

0/25 multipore $\rightarrow$ $\frac{1}{5}$ 0/3 multipore

in sack + 1000 μ l ($\sim 1 \times 10^6$ cpm.) of Harveys
HF 11-49 cDNA probe + 2 ml Denhardt's Hybridizing
~~Hybridizing~~ Sol. + 1 ml ~~sol~~ 0.57 2X Hybridizing

To 70°C at 1.30 pm

Save this probe.
Recovered 3.5 ml

11.45 am to 70°C with 2XSSC + 2% SDS.

↓ Fri, after ride

12.45 pm $\rightarrow$ 1.40 pm $\rightarrow$ 2.40 pm

Rinsed 2XSSC. Dried. To autovar 3 pm
Off


Unfortunately multipore stuck to
film (not quite by)
Again Sat 2 pm $\rightarrow$ Mon 9 am

11. am
Scl
0/25

Mon $\rightarrow$ Sat noon

Good result

Important (technical comment)



No stable
transfer by
hydro - with 48 hrs
antioxid of label


Thurs. March 24th

Small piece gels

Fred suggested electrophoresis out
out acrylamide might work. Meanwhile
try high conc. agarose.

Fri, March 25th

Tried an old Haettgel (5% acryl) of
LAF with ~ 15 min SEC: I ~ 15 min SEC: II
 ~ 15 min ep. at ≥ 1 v/cm in 6xSSC (15 Amps ~ 2 v/cm
in old testainer) Active J



Rinsed. To
Decharth's at 4.30 pm
↓ 70°
11.0 am

Remained excess replaced
by used probe P+1 & P-1 from 081
to 70° Sat. am

Sun Noon Probe recovered
Rinsed 2x to 70° w/ah
2xSSC $\frac{1}{2}$ % SDS
3 pm → 4.30 pm

0133 USED
PROBE

2xSSC rinse. Auto. 5 pm

Mon 10 am → Wed 10 am Negative Mon 9 am


Result shows transfer occurred on
both attempts 15 min $\rightarrow$ 45 min. so if
geometry is improved, the method will work!



Mon. March 28th

More serious gel transfer expt.

LAF gel 1135 containing $\sim 60,000$ cpm
(of which $\sim 30,000$ is in Hha Δ lac piece
 $\rightarrow$ rest in undigested Δ lac).
in outside slot (of 6).



V/cm sh'd be $> 1V/cm$

4.45 pm Counts still in gel

5.05 pm there again (5.35 pm pH 9.7)

$\downarrow$
5.50 pm

Most of counts still in start
gel - must channel current
better - none appear to
be through - same on Multipore

fixed $\rightarrow$
to control
6 pm $\rightarrow$ 8 pm

0137

FRONT POST TRANSFER

POST TRANSFER

Mon March 28th

Hae III / Bam again

On 9:35 pm

Air cooled.

LAF DB1	Hb4	Hae III	5%	Agarose (autoclaved)
LAF DB1	Hb4	Hae III / Bam		
LAF DB1	Δ lac	Hae III / Bam	4x H	
LAF DB1	Δ lac	Hae III	2.5 V/cm	100 mA

At 11:50 pm $\searrow$ to 1.0 V/cm.

Off 8:30 am (slots had dried & current was ~0 mA)

Photographed. Sol. I 8:35 am $\rightarrow$ 9.0 Sol. II 9.0 $\rightarrow$ 9.25 am

(Top down)

Transferred as usual from 9:30 to 4 pm
 Stained & checked. Transfer incomplete
 Denhardt at 70°. Use overnight in future

Hybridized 11.0 am tiled $\rightarrow$ Thurs 8:30 am
 against 0/31 (3.5 ml) probe

2xSSC $\frac{1}{2}$ 9 am $\rightarrow$ 10:15 am 10:20 am $\rightarrow$ 2
 70°C $\rightarrow$ Rinse

Negative

2 $\rightarrow$ 4:30 2xSSC Rinse & dry. Autorad 5 pm
 Sat noon $\leftarrow$ ② Fri 10 am $\leftarrow$ ①


Wed - April 30th

E.p. transfer

Rough 5% gel run with $\sim 30,000$ cpm

12 noon
~~12 noon~~

Take piece from #63
to Soln. I 12.20 Gen II



$\text{pH}_0 \sim 8.5$
12.55 pm On to ep pH in 20X SSC 15A ~ 6 v across no path
around gel (2-3 v/cm)

Fruity smell
not much in

$\text{pH}_0 \sim 8.5$	$\text{pH}_{45\text{min}} 8.5$ (getting warm)
$\text{pH}_{10\text{min}} 8.9$	$\text{pH}_{70\text{min}} 8.5$ (56°C)
$\text{pH}_{20\text{min}} 8.7$	$\text{pH}_{100} 8.35$ (62°C)
	$\text{pH}_{180} 8.4$ (65°C) $\text{Gel} \sim 2 \text{ v/cm}$

No counts on Mullipore. Appearance


Check after 2 hrs. / suggests gel may have stopped below mask. Counts still in gel. Cleaned up apparatus.

Replaced 20xSSC.

Redenatured DNA in sample

Reset up without mask

— should not alter V/cm anyway.



5.15pm 4V total obtainable with 15A without mask.



9.30pm Off. Recovered in morning. Can't find counts on Mullipore — but they fit in leave gel.

This is all satisfactory — counts are, in any case, too near origin. Repeat.

10.30th am
Autorad

→ Sat 12 noon


Negative

Thurs. March 31st

Lesley's latest gel ep

Secⁿ I ~15 min Secⁿ II ~15 min

Gpm ~4 v/cm



12 midnight

Still no counts moving! Crazy

Antorad 12:20 am → 10 am

Nothing!

Negative

Donna's Marker mix

495 base pairs = 192x

2.6

3.7

5.4

6.8 (nt. end)

9.1

9.8

11.8

12.2

14.0

15.4

(17.7) partial see a very high conc.

21.3

(25.8) partial " " "

44

51 fusion (44+6.8)

60 441 Refund

(65) see if had sample fusion (44+21)


(67) " " " (60+6.8)

fusion (60+21.3)

80

100



? absent



146

1265 014702 0147 146 days

~8 days



Result with λ probe
shows two faint
bands at 12 hr. ~~0147~~
~~0147~~ - got two
pieces muddled

Wes
+ Monish Set up the other half  with H₂A probe
0127 200 μ l (now $\approx$ 750,000 cpm) in 200 μ l 2x
annealing $\rightarrow$ 2 ml 1x

0147 3pm

Thurs 3pm 2xSSC $\frac{1}{2}$ SDS $\rightarrow$ 4.15 $\rightarrow$ 5.30 $\rightarrow$ 8.0

0146 Autoraf. 8.30pm $\rightarrow$ 8.30am



slots

Two sets
of two bands

Fri
April 8

Sat
April 16



See back
on 145

Did not work
- too much loss.

147

Sunday April 3rd

Background test

Sat April 16th
o146 piece soaked
in 0.4M Na_2SO_3
0.2M K_2CO_3
NaOH to pH 11.9
for ~5 min. Rinsed
in 2XSSC & dried
Automat.

Sat
16th

Sat
30th

100 μl of 45 Bulk Mouse R1 digest (last of
to remainder (~10 μl) of 51 10X R1
to give

~ 110 μl o147 2X Mouse R1

Sat April 16th

o147
~ 55 μl
Sams
12 hrs

55 μl + 1 μl o145 Marker mix

On 12.20pm Off 4pm

o147 2X Mouse R1	0.7% Agarose
o147 2X Mouse R1 + λ	4X H
o147 2X Mouse R1	~2.5 V/cm
o147 2X Mouse R1 + λ	100mA
o147 2X Mouse R1	
o147 2X Mouse R1 + λ	

Auto 8pm
(doesn't look
good)

5.30

2X

Ex

80

3.10pm

1/2SSS
2XSSC

2pm

Stained. Photographed. 15 min Solⁿ I 15 min Solⁿ II

Transferred as usual

5pm



9am

80° under vac. 10.10am → 12.30

~ 12.30pm → In Denhart's at 70° 10.30pm

1/2 off & plus o133 probe Rest back to Denhart's
- both rather little 2pm freeze

Apr 16th

Grunstein - Hogness

9cm Millipore .45 μ

boil 3 x 1 min in H_2O in petri dish

autoclave between blotting 10' dry in autoclave 10'

place millipore on ~~petri~~ ^{agar} medium in petri dish no bubbles
replica plate onto millipore filter and grow colonies

The Procedure

- 1) Place filter gently on a blotter moist with .5N NaOH 7 min no air bubbles
- 2) Place filter gently on a blotter moist with 1M Tris 7.5 2 min
- 3) repeat step 2
- 4) Place filter on a blotter moist with .5M Tris 7.5 1.5M NaCl 5-10 min
- 5) Place filter on filter separates and dry with hard vacuum 2 min

optional → 6) ~~2mg/ml~~ Soak filter in 2mg/ml ^{predigested 2 hrs 51°C} Pronase 1XSSC
15 min dry with vacuum on apparatus

7) Wash filter on filter apparatus with ~100ml 95% EtOH

optional → 8) Wash filter or filter apparatus with ~100ml chloroform

9) Dry with vacuum on apparatus

10) Wash filter with .3M NaCl on vacuum apparatus

11) Dry 80° in vacuo 3 hrs to overnight
and handle as any microcellulose paper
hybridization

We hybridize in

chlorinated by 2 hrs
stirring in amberlite
50% formamide and filter
4 x SSC
20 mM Tris 8.3
15% SDS

Normally 37°C (rDNA)

for fibron mRNA 50°C

Granstein - Hogness

9cm Millipore .45 μ

boil 3 x 1min in H_2O in petri dish

autoclave between blotting 10' dry in autoclave 10'

place millipore on ~~petri~~ ^{agar} medium in petri dish no bubbles
replica plate onto millipore filter and grow colonies

The Procedure

- 1) Place filter gently on a blotter moist with
.5N NaOH 7min no air bubbles
- 2) Place filter gently on a blotter moist with
1M Tris 7.5 2min
- 3) repeat step 2
- 4) Place filter on a blotter moist with
.5M Tris 7.5 1.5M NaCl 5-10min
- 5) Place filter on filter apparatus and dry with
hard vacuum 2min

K111

5/5 Prestidge modification of the Grunstein filter screening procedure

- (1) With toothpicks stab the colonies to be tested in an ordered array on an L plate (the master plate) and grow overnight at 37°C. *A stainless steel template up to 150 colonies is helpful.*
- (2) Replicate the master to a nitrocellulose filter on an L plate and grow overnight at 37°C. The filter will remain on the surface of the agar while you replicate and the cells will grow normally on the filter.

Store the master plate at 4°C.

We punch our own filters from 293mm diameter HAWP Millipore filters. These are boiled three times in water to remove detergent and then autoclaved (layered between filter paper) for 5 min. (fast exhaust) to sterilise them. The filters shrink a large and variable amount under prolonged autoclaving so it is important to keep to this time. Millipore supply a 90mm HAWP filter which is too large to fit into a standard petri dish (88mm internal diameter). I have been told that these shrink sufficiently during autoclaving to fit into a petri dish but we have not tried this.

The following steps are performed by placing the filter on blotting paper saturated with the appropriate salts.

- (3) Place filter on 0.5N NaOH for 7min.
- (4) Transfer to 1.0M Tris/HCl pH 7.4 for 2min.
- (5) Transfer to a second 1.0M Tris/HCl pH 7.4 for 2min.
- (6) Transfer to 1.5M NaCl, 0.5M Tris/HCl pH 7.4 for 4min.
- (7) Transfer to a filter block and suck hard for 1min.
- (8) Wash with 2x30ml. EtOH and suck dry.
- (9) Bake under vacuum at 80°C for 2hrs.
- (10) Moisten the filter with the appropriate ³²P labelled RNA in 50% formamide 5xssc. and incubate under oil for about 12 hrs. at 42°C. It takes about 0.7ml to moisten a filter.

We use >5x10⁵ cpm. per filter where possible. For a discussion of this hybridisation step see Grunstein and Hogness PNAS in press.

- (11) Wash the filter as follows:

2x30min. in 50% formamide 5xssc. at 42°C
2x30min. in 2xssc. at room temperature
1x30min. in 10µg./ml. RNase in 2xssc. at RT
2x30min. in 2xssc. at RT

30% 8.5x SSC
- 2M PB

5x SSC

5x SSC

deal for pBR 322

Pat

May 20, 1976

Pat,

I'll try to describe the procedure completely. We don't have a protocol annotations written up yet, but I'll send you one when we do. We're growing with $\phi\phi$ bugs growing in individual wells in Microtiter plates so our reagents are different than yours. To grow up colonies on filters for hybridization, we use cookie sheets with fitted aluminum lids. These are lined with hard agar with tetracycline and 6 $4\frac{1}{2}$ " squares of Whatman 3MM are placed on the surface and allowed to wet thru. Then we replicate a 12x12 array of bugs on to the surface of the 3MM squares with a 144 "bed of nails" replicator. The bugs are grown overnight at 37°. Bacteria growing over tetracycline containing medium become intensely fluorescent so growth is readily assayed by looking at the filters with UV.

(By the way to prepare the filters we simply cut $4\frac{1}{2}$ " squares from 3MM paper and autoclave them. You can number them with a Sharpie marking before autoclaving and the label will survive all the subsequent procedures. Overgrowth of bacteria immediately over the marking is inhibited. If you are too near the edge, you run the risk of losing the marking due to accidental amputation during batch workup of the filters, so you may want to check each filter twice if you do this.) To work up the filters we simply plop into a plastic dish pan with 0.5 N NaOH and wash them gently for about 10 minutes and then change the old stuff with fresh 0.5 N NaOH for another 10 minutes. Then we wash the filters twice with 0.5 M Tris for 2.5 minutes

Then we wash twice with 1 x SSC, again for 2.5 minutes each. Then we wash twice with 95% ethanol for 2.5 minutes each and allow the filters to dry. Hybridization is done at 37° overnight or longer in 50% Formamide. We try to have 100,000 CPM/ml of pure probe. Then autoradiography a few days gives black spots. After hybridization we wash the filter 3 times with 2XSSC and then RNase at 37° for 1 hr with 20 micrograms/ml pancreatic RNase. If high background is a real problem, you can wash the filters in Waltham tap water or 0.01 x SSC. This reduces the level of hybridization somewhat. But we really haven't had problems with background, usually just wash the filters with 2 X SSC. It frequently takes a lot of washing; we usually monitor the supernatant by Cherenkov counting. The filters are usually still hot by monitor (Geiger tube) (not very hot), give lovely radioautograms. Be gentle with the filters, they're only fragile. If you see hunks of filters floating around during the washes, slow

Air dry when done washing and autoradiograph.

Miscellaneous: It takes about 6 ml to wet our $4\frac{1}{2}$ " squares of 3 MM. We store them in plastic sandwich boxes or taller freezer containers. The lab number is 617-647-2434 if you have any questions. I'm also sending a latest poop sheet from Stanford on techniques for nitrocellulose filters: we've simplified denaturation and washing of the filters by simply placing filters on paper soaked with the different reagents. Igor introduced us to the use of paper. He noticed that when colonies are lost from the nitrocellulose during washing, there wasn't hybridization. Reasoning that a "DNA print" was not the source of hybridization, he tried paper filters since the bugs grow down into the paper and thus not be lost during processing.

Sincerely,

Ralph

Will be out of town May 21 - 28

pH 7.4

Monday - April 4th

Hogness re-visited

70mm filter papers available from
stores are: -

1, 2, 40, 42 & 50

~10 of each washed on Boudier with
hot S.D.W. Autoclaved.

Tues. April 5th

Decided to test with a λ probe on 4 ml ($\pm \lambda$)
- this is more stringent for #'s of copies but
less since λ is so big.

Sat. cultures available - assume $\sim 1 \times 10^9$ / ml
- took 1 ml of ~~KH802~~ ^{3/29} + 0.1 ml ~~KH802~~ ^(4/4)
Diluted to 10^{-5} (i.e. $\sim 10^4$ / ml) $\rightarrow 10^{-6}$ ($\sim 10^3$ / ml)

Spread 0.1 ml on the papers over
NZ plates - papers pre-wet in NZ


- #1 | more fragile
- #2 | - too full of liquid
- #40 | Tougher
- #42 | - nice water content
- #50 | too unstable but
- but better than
1 & 2

longer sit before spreading
 10^{-5} gave same conclusion.

To 37°

3:45 pm

12 pm
Wed. growing OK



Wed pm looks as if the thinner papers give bigger colonies more quickly.



Thurs am — heavy per filter paper growth
Stained colonies. Clear that radial transport
(AB, dilute, in 50/50/10 has occurred — presumably
with uptake) by evaporation.

Repeat & seal plates — also put
papers down



Fri am Checked re-incubation of stopped
dishes — clear that bacteria are through
OK but removal smears them
e.g. X1776 #42 5.1

X1776 #40 5.1



So will have to pick colonies
or print a replica. Try that on next batch.