

Jan. 16th 1994.

Smith
Dept. of Pathology
University of North Carolina
Chapel Hill
NC 27599-7525

The Boorum & Pease® Quality Guarantee

The materials and craftsmanship that went into this product are of the finest quality. The pages are thread sewn, meaning they're bound to stay bound. The inks are moisture resistant and will not smear. And the uniform quality of the paper assures consistent rulings, excellent writing surface and erasability. If, at any time during normal use, this product does not perform to your expectations, we will replace it free of charge. Simply write to us:

Boorum & Pease Company

71 Clinton Road, Garden City, NY 11530

Attn: Marketing Services

Any correspondence should include the code number printed at the bottom of this page as well as the book title stamped at the bottom of the spine.

One Good Book Deserves Many Others.

Look for the complete line of Boorum & Pease® Columnar, Journal, and Record books. Custom-designed books also available by special order. For more information about our Customized Book Program, contact your office products dealer. See back cover for other books in this series.


Made in U.S.A.

RMI290393


Sun. Jan 16th

ANTRA: DUP (cont.)


Proposed constructs for dupⁿ or delⁿ



The deletion will require 5' flank with opposite site; the 3' flank opposite is already made.



Checks on 5' flank
or 3' flank
for Xho on next page
→



Dye
Water
B'3 3'C
B'3 3'X
B'3 3'H
W
B'3 5'C
B'3 5'X
W
Marker
W
Dye

Conclusions: No Xho I site in 1.3. 3' flank.
 2 Xho sites in ϕ 2B2, but not in Paula's
 candidate to 5' flank. Proceed.
 No Xho I ^{sites} in Charon 35 vector

CAUTION This ^{may} not be a rigorous test since we
 have to assume the 9.5 kb ~~could~~ give $6.5 \text{ kb} + 2.8 + \sim 0.6$
 But the 6.5 could have given $2 \times 2.8 + 0.6$ - unlikely
 Anyway test again when fragment
 available.

0.88 HX4 MAX
On 2:36pm
Off 3:36pm

3

Sun. Jan 16 d

Checks on 5' & 3' flanks for Xho sites





} 3' flank

Take ✓ 6 µl 10X R1 salt

✓ 50 µl ddw

✓ 4.5 µl A14.8 #26 mini 3' flank

} 5' flank



Test

37° 12:25 → 1:25 pm

Take ✓ 4 µl 10X R1 salt


✓ 26 µl ddw

✓ 10 µl Paula's Charon 35 clone 2B2



✓ 2 µl Eag1 (20u) lot 21

✓ 1 µl Sal1 (20u) lot 47

SHS 66
2 B2
0.1 pg/µl



Test



07183 5' Oligo Name: OSEAGXHO

made: 1-16-94

G	G	C	C	C	C	T	C	G	A	G	G	

FT11: 12


C: 83%

500 Å Prime Synthesis 004960


Back up into mlTE.1XCIAA
Stored as B'5 OSEAGXHO

Sun. Jan 16th


Eag I to Xho I



Xho I
8 bp overlap. Eag removed. Xho I inserted.



Conclusion: looked like
neither cut at all!
Prepared fast band
(roughly) anyway. Good
cut for 148 Blue
-moderate for 155 Red (Black)



Mon. Jan 17th

3' flank preparations

Sylvia made the two orientations in supermini as

A'148 H13X and A'155 29 X/H
 O.S. → (#26) O.S. → (#29)

The vector is ori-vector, ~2 kb, so can separate ok.

Take ✓ 21 µl (3 µg) A'148
 ✓ 3 µl 10X R1 salts
 ✓ 4 µl ddw
 ✓ 1 µl (20v) Hind 3 lot 45
 ✓ 1 µl (20v) Xba 1 lot 29

Blue
 B'17
 148 X/H
 0.1 µg/µl

o ✓ 14 µl (3 µg) A'155
 ✓ 3 µl 10X R1 salts
 ✓ 11 µl ddw
 ✓ 1 µl (20v) Hind 3 lot 45
 ✓ 1 µl (20v) Xba 1 lot 29

Red
 B'17
 155 X/H
 0.1 µg/µl


37° 4:15 pm to 5:45 pm

8:05 pm to 520 mA


On 6:35 pm Off 10:15 pm

1.5 V/cm 3 Max 260 mA

(~0.25 µg/sample lane)




ori vector
 1.3 kb
 ssDNA



↑
 single cut

Conclusion: Dilution works
 - or gel is better??

Reaction went to completion,
 probably, with rather
 a lot of ssDNA.
 There is not single cut
 plasmid - it migrates
 at 3.3 kb



Conclusion:

Must have been
 gel resolution
 all along. Repeat
 digest & try
 using 1.2% agarose. (HX)
 or test sample first.
 3'9 37°

10:35 pm → 9.15a

Tues. Jan. 18th

X 11.3 H digestion check

40pl Excess of B'7 148 X/H diluted + 4pl 10X R1 salts

20pl	20pl
W	W
X/H dil ^d	X/H dil ^d
X/H ² dil ^d	X/H ² dil ^d
W	W
Marker	Marker
W	W
Dye	Dye

37° 8.55 am to 10.50 am

On 10:47 am off 11.30 am
0.8% H x 4 max

(Blue)

(Red)

(B'9²)

Take 21pl (3pl) A'148
X 18pl 10X R1 salts
X 139pl ddw

X 14pl (3pl) A'155
X 18pl
X 146pl ddw

A'9
148 X/H

X 1 pl (20u) Hinf 3 lot 45
X 1 pl (20u) Xba 1 lot 29

X 1 pl Hinf
X 1 pl Xba

A'9
155 X/H

Dye water

(Blue) 148
~~water~~
~~dye~~
~~water~~

37° 12:30 pm to 2.45 pm

(Red) 155
~~water~~
~~dye~~
~~water~~

On 4:10 pm

Pause 2.45 pm - 9.0 pm
Off pm


2.5
500 am A' V/am

should be ~ 10 pm

Blues run until  saplogue



Conclusion: Run until
Blue 2 is ~6cm
Cuts are fine (less of 55)
SS DNA now
well out of way.



Conclusion: gel
over-run (fure alarm!!)
but cuts pretty good.
Proceed ~~with~~ 
+ 10% 8M NHE₄Ac
Melted 80-85



Wed. Jan 19th

H 1.3 X prep. (carb.)

Xba 1.3 Hnd
Hnd 1.3 Xba

On 10.02am Off 11.15am

Dye
W

~1.7% Soap. HX1

B'9² 148 X/H² Blue

W

6 V/cm 100mA

B'9² 155 X/H² Red

W

Marker

Run until yellow dye is at wick
(in this case 80 mm)

W

Dye

On 12:40 pm

Off 3:40 pm

~1.7% Soap

6 V/cm 210mA

2nd blue almost at wick

Dye

W

148 X/H
B'9²

Blue

same left

B'11 148 Blue

W

Dye

W

155 X/H
B'9²

Red

None left

B'11 155 Red

W

W

Dye

* ~1 ml 40% H₂O₂
Cent. 1 B'11 148 Blue
2X CIAA B'11 155 Red
Total vols. ~1.3-1.4 ml

Tools

130 pl of each ✓
+ 390 pl EtOH ✓

2 back into 15 pl
for test rep at

B'11 Core 155 Red

B'11 Core 148 Blue

Gel (over) →



Conclusion: Both are fine -
a) expected (48 is at
highest conc.

Wed. Jan 19th

Subcloning Sal 5' flank Eag

Cannot conveniently change Eag to Xho & use
blue script since Sal & Xho are too close.

∴ subclone into BS & then change site. Paula needs
this route anyway. Let her do it.

later thought (Thurs) w131 pSP64 Sac → Xho can be
used - it has a linker! third Pst (Sal) Acclinc? Kba Bam Avc Sma (M)

Thurs. Jan 20th

On 10:58am

Tests of II.3 fragment

Off 11:35am

0.8% Hx4
max

	Dye
	w
~120pl	Blue B' 11 148 conc $\approx$ 120pl unconcentrated
"	Red B' 11 155 conc $\approx$ 120pl "
	w
	Marker
	w
	Dye

On 11-19 am
Off 12-15 pm



Conclusion:

⊕ Egg digest

is fine.

Sal used ~~not~~ of Karmid used

not complete (only 40%)

on 2A

continue digest

- see

B15

Sal/Kho² below

Dye
W

Post Egg

Post Sal

Post Sal

Post Xho

Post Xho

W

Marker

W




Dye

0.86 Hx4
max

Thurs. Jan 20th

Salt 5' Blank ~~Eag~~ → Xho2B2 ϕ Xho
Sal
2A
plannedTake 3 ϕ 10x RI salts✓ 15 ϕ ddw✓ 10 ϕ (2 μ g) Paula's 2B2 B'3 B'15
2B2/Eag✓ 2 ϕ Eag 1 lot 21 (200 total) 0.01 μ g/ ϕ 37° 5.07 μ m to 7.30 μ m (frozen) 2+15 B'15
Post EagTake 3.2 ϕ (4 μ g) W131 #2A SHR 26
✓ +6 ϕ 10x RI salts 1.24 μ g/ ϕ ✓ 30 ϕ 51 ϕ ddw+ ✓ 1/2 ϕ (100) Sal I Lot 4737° 5.07 μ m to 5.52 μ mB'15
2A Sal0.04 μ g/ ϕ + 1/2 ϕ (100) Lot 26+ 1/2 ϕ (100) Xho I Lot 262+15 Post
Sal Frozen2+15 Frozen
Post
Xho6.00 μ m to 7.30 μ m (frozen)2+15 Post
Sal/Xho+ 1/2 ϕ (100) Sal I Lot 472+15 Post
Xho/SalFin. Pooled the Sal/Xho 2A
+ Xho/Sal 2A. Added equal
vol. IXRI 2 added 1 ϕ more of Sal
Xho slack to 37° 12.55 μ m to 3.45 μ m
asB'15
Sal/Xho²~ 0.04 μ g/ ϕ

(contd.) Two gel slices processed as usual 10% SMNH_4Ac 75-85°C
 PCH/HOQ, 2x CIAA.



B'17
 Sal 5' flank Kho
 Cuts at 35 mm
 or 31 mm

Conclusion ①:
 Missed partly!
 Recovered stained
 lower rest for
 B'17 probe
 - can use it also
 to check Kho

② Conclusion: Repeat prepⁿ, but
 continue with this 5' flank fragment

Tr. Jan 21st

Sel 5' flank Egg $\rightarrow$ Xho (could.)

Take whole of B'15 2B2/Eag

- ✓ + 6 μ l (est^d 3 μ g) B'15 OSEAG-XHO
- ✓ + 4 μ l 10 $\times$ p95 buffer
- ✓ + 4 μ l TA ligase

B'15
2B2 Eag
ligⁿ

15 $^{\circ}$ C 10:40 am to 11:25 am 2/3/86!

R.T. 11:25 am to 12:05 pm

37 $^{\circ}$ 12:05 pm to 1:15 pm

2 μ l + 15 μ l
B'15 part
ligⁿ

40 μ l Blue 5' FLANK
45 μ l Red 5' FLANK
PROBE CONC
W
Marker
W
Dye

+ 4 μ l 8M NH₄Ac

+ 132 μ l EtOH

Back up into

5 μ l LdW
10 $\times$ RS + 150 mM NaO
6 μ l ~~XXXXXX~~
1 μ l Sel 1 (20 ν) lot

B'17
2B2
Eag/Sal
Xho

37 $^{\circ}$ 2:07 pm to 3:47 pm
Diluted with 30 μ l LdW
o applied to gel

$\approx$ 0.05 μ g/ μ l

① Dye
W
1
2
3
4
5
6
W
Dye

0.8%
Hx4
Seap

0.8%
Agarose
Hx6

Not quite as much sample



On 4:10 pm
Off 9:10 pm
 $\approx$ 3V/cm
450 mA

Mislabelled
as B'18 over page
←

Sunday, Jan. 23rd

ANPRADUP lig.

All fragments now available with ambiguity in 3' flank. ^{but one} assemble two ligations (Red & Blue) & proceed. (Error) Sal I ~~Neo~~ hint, still to make.



Take (7.7 μ l (10 μ g) w/ 125 #9 SHR34 VPCR 1.30 μ g/ μ l B'19

10 μ l 10x R1 salts

82 μ l ddw

1.5 μ l Sal I (30 u) lot 47

37° 2:25 pm to 5:50 pm

+1.5 μ l (30 u) Hind III lot 45

37° 2:25 to 5:50 pm

2 μ l + 15 μ l

2 μ l + 15 μ l
B'19
Pre

Use this only for probe


Repeat over

B'19
Neo
Sal/Hind

2 μ l + 15 μ l
B'19
Post
Sal


B'19
Post
Sal/Hind

①



22 mm
29 mm cut


①



① Conclusion: Adequate (just) cut.
Proceed. $870 \text{ mg} + 90 \mu\text{L SM}$
 $80^\circ \text{ Fk } \phi\text{OH/HOQ}$

B'20
pSP64 Sal/Xho $2 \times \text{CIAA}$

②



← Post Sal

← Post Sal/Hind

← Post Sal/Hind

Total vol $\approx 800 \mu\text{L}$

② Conclusion

The digestion was
Good: Sal OK, Hind OK.

Cuts at 40 mm

35 mm



30 mm

25 mm

Processed to $\phi\text{OH/HOQ}$
 $2 \times \text{CIAA}$.


75 μL of each plus 210 μL EtOH
* back up into 15 μL for gel test (B'22)

B'20
Hind/Neo/Sal
M


Mon. Jan 20th

Can use dye, since it is only vector, but don't need it.

Take 50 μ l B'15 Sal/Xho 2 + 50 μ l ddw. ~~load~~



①



on 10.35am 0.8%
off 11.45am Hx4
max

①

Dye
w

2+13

Pre B'23 high

Post B'23 high

75pl

B'20 F

B'20 M

5X

B'20 S

w

Marker

w

Dye

① Conclusion: B'20 M is good
- save.
high: worked -
↓ vector & ↑ fragt
on repeats.

7.5

② See over

②



last trip

DH5α 1/7

~180pe

~35pe Rest of

B'23 high

0°

11.26am to 11.58

42°

90 sec

Plated 37° 12.15pm

10am

14 colonies

or "halos"

Mines attempted

10.25am Sat 5.25pm

↓ stationary 37°

② Conclusion: Not this time!

Mon. Jan 24th

Subcloning of Sal [5' flank] Xho in pSP64

Rough attempt without gels.

Take ✓ 210 μ l 5' FLANK B'18 FRAG (blue)

✓ + 40 μ l B'20 pSP64 Sal/Xho

✓ + 150 μ l EtOH

Back up into 10 μ l 1x p95 lig⁺ buffer

B'23
pSP5' FLANK
lig⁺

2+13
PRE
B'23

15° 5:11 pm to 6:04 pm

R.T 6:05 pm to 6:35 pm

37° 6:35 pm to 6:45 pm

R.T 6:45 pm to 8:30 pm

cold room
overnight

2+13

POST
B'23

Transformation

B'23 pSP64/5' FLANK

B'23

DH5 α 12/6

80 μ l ✓

~~25 μ l~~
DH5 α
B'23 O.N.

40 μ l

B'23 lig⁺

2 μ l ✓

2 μ l

0°

8:35 pm to 9:18 pm

11 am 0°

~ 12 pm

42°

90 sec

2° ~ 90 sec

Plated

Rest 20, 5, 20, 5

37° 9:30 pm

Ther
12:15 pm
37° 12:15 pm to 1:44 pm

9:30 am looks like 0
Repeat with better cells.

Catch up gel

Sun Jan 30th

24

Sun Jan 30th

Dye

W

Transformation

B²L¹

Blue

B²L²

Red

B²L¹

Blue

B²L²

Red

Marker

W

2 #13

Pre B²5B

DH5 α 1/7

~100 pl

~80 pl

120 pl

120 pl

Post B²5B

high

1 pl
B²5 Blue
high

1 pl
B²5 Red
high

1 pl
Blue
high

1 pl
Red
high

Pre B²5A

Post B²5A

W

0.8% HX4

max

~11 am

to ~12 pm

0° 11.01 am to 11.34 am ~5.29 pm

90 sec 42° ✓

Plated Rest

Red B²L¹

20, 5, 20 / 16, 5 / 16

Blue

11.54 am Sun

B²L¹

B²2
minis
←
See
over

B²2
min
← see over



Conclusion: Quite good proportions;
didn't need quite as much B²1/155 Red
ligations are promising.

Size sh'd be ~14 kb

1.0	Neo
1.3	3' flank
4.7	TK Vector
6.4	5' flank
13.5	

DMS
1/7

pl
Red
h^g

N 5.20 pm

4

She

2. 10

(29)

;

5 Red

1200 total
3 Flank
Blue


Important
F10ys
Have fun

Tri. Jan 28th

First attempt at making complete construct

for NPR Application (see §19)

Trigments are now available:-



B25 Blue high

B25 Red ligⁿ

Take ✓ 20 pl A'γ Xho TK Xba M

60 ml B'20 Hand Neo Sol M

(L) 230 pl B'18 5' FLANK FRAG (Blue) whole sample

100 100

B'U / 158 B...

STA# (3 vol)

+80 ✓

✓
10pl 1x p95

9.10 + 13.4 | PBE B'15

+ 1 μ l T4 ligase

15° - 52 mm x 1:28 mm

27 0:29 to 6:33

00221000

10 pl 1 x p95

$$P \cup P' = P \cup P' = P \cup P'$$
$$+ \uparrow \text{NP Th. } \text{div. } \text{no.}$$

Too cold - 10:15 am Sun

2 pet 13 POST Blue R25

9.12 + 13 POST Red B25

Nathropinetic peptide receptor 1
(old name ANPRA)

25

Oligo # 07236

Date made: _____

LENGTH: 12% G+C: 50%

5'

note: OS ECOBAM

A A T

T G G

G A T

C C C

[] [] []

[] [] []

[] [] []

[] [] []

500 Å
0.2 uMNAC-020-1
Prime Synt
0 014

5'

note: _____

10:01:49A , 1/29/94

Run ID

- 394 Synthesis Setup Listing -

(Version 2.0)

Column 1

1:53:40P , 1/29/94

Run ID :

Cycle : 02UM USER

End Proc: End DESS (DMT = Off)

Sequence: 07236

Total bases = 12


A= 3, G= 3, C= 3, T= 3, S= 0, 6= 0, 7= 0, 8= 0
(mixed bases= 0)

MW: 3643.4

5' > AAT TGG GAT CCC <3'

Sat. Jan 29th

Eco RI $\rightarrow$ Bam HI





Back up into 500 μ l of TE 1X CIAA

2 stored as B26 OSECOBAM

B'24
8 Blue

B'24
26 Red

streaked out from Wed.



Conclusion: Missed! Xho not there

Xba linearized to ~~9.2 kb~~ ~ 9 kb

Xba/Hind → 1.3 kb + 4 kb

Hind/Sal → 4.0 + 3.2

~~approximately 1.8 kb~~

So it has 1 Xba site


2 Hind sites

1 Sal site

just try some more!

Design Error

Sal will ligate into Xho



on 3:10 pm
 4:40 pm
 0.86 HX4
 WAN

Wed. Feb 2nd

1st clonings of ~~APR~~ DUP

Sylvia picked the very dispersed, satellite-rich, columnar streaked out to make mini. Some labelling problem of gels but clear result

6/9 Blue look good; 18/25 Red look good.

Proceed with 8 Blue and 26 Red

KOAc sups. + 350 μ l isopropanol.

Ppts. washed. Back into 50 μ l COIM-OTA or

B'24
8 Blue

B'24
26 Red

Unaut

Xho

X/Sal

X/Xba

X/Xba/Hind

X/Hind/Sal

W

Unaut

Xho

X/Sal

X/Xba

X/Xba/Hind

X/Hind/Sal

W

Marker

W

Dye

Take 32 μ l 10X R1 salts

+ 272 μ l ddw

+ 16 μ l B'24 8 Blue

+ 4 μ l (5X160) Xho1 lot 26

32 μ l 10X R1 salts

+ 272 μ l ddw

+ 16 μ l B'24 26 Red

+ 4 μ l Xho1

8 Blue
Xho

8 Blue
X/Sal

8 Blue
X/Xba

8 Blue
X/Xba/Hind

8 Blue
X/Hind/Sal

26 Red
Xho

26 Red
X/Sal

26 Red
X/Xba

26 Red
X/Xba/Hind

26 Red
X/Hind/Sal

32° 12:15 pm

2:30 pm

30

on 2:02 pm

(3)

Dye
Mark

55

56

57

58

59

60

61

62

63

64

65

66

Wat

67

68

69

70

71

72

73

74

75

76


77

78

79

80

Dye



② on 12:50 pm

Dye

W

25

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

Marker

Dye

W

Dye

on 11:42 am

Dye

W

1

2

3

4

5

6

7

8

9

10

11

12

13

14

Marker

Marker

15

16

17

18

19

20

21

22

23

24

W

Dye

Thurs. Feb 3rd

B¹24² minis

Picked early & still not v. satellite except #6 & #7, 14 & B¹24² BLUE colonies (#14 not streaked). To 37° at 10:25am

B¹24² RED #15-25

and #31-55 (no 26-30)

RED

→ 37° 11:25am

B¹24²
Blue


RED #56 → #80 (only ~6 more left)

37° 11:55am

- ① Conclusion: None of Blues are correct or 15-24 of Reds. Finish the other two gels. If any Red is correct, repeat Blue. If none - proceed with correction strategy. Probably these minis were a waste of effort because concⁿ of 5' flank Sal-Xho is low & is ↓ to 50% by orientation AND can self ligate.
- ② & ③ - confirm the problem

Glorious yield of the 3 fragment construct: probably closer to 7.1kb than 9

B¹24²
Red



3

Dye
Max

55

56

57

58

59

60

61

62

63

64

65

66

W

6

6

Into 500 μ l TE 1X CIAA
Assume 0.5 μ g/ μ l

B'32 OS
Xho I
Not I Xho I

07247

de:2-3-94

II: 12

: 83%

5'

note: OS Xho Not Xho

T	C	G	A	G	C	G	G	C	C	G	C	

Run ID :

Cycle : 02UM USER

End Proc: End CESS (DMT = Off)

Sequence: 07247

Total bases = 12


A= 1, G= 5, C= 5, T= 1, S= 0, 6= 0, 7= 0, 8= 0
(mixed bases= 0)

MW: 3643.4

5'> TCG AGC GGC CGC <3'

Thurs. Feb 2nd 1990


Correction of NPR I DVP design




Requires changing

(There are
2 Eag I
sites in
pMCITK
but no Not I.

Two have same overlap)



OS Xho Not Xho $\equiv$ Xho Eag Xho




(contd. $\rightarrow$)

Nucleic Acids

pSP64 Vector

Product	Size	Cat. #
pSP64 Vector	20µg	P1091



GGCTG CAGGT CGACT CTAGA GGATC CCCGG GCGAG CTCGA ATTC...3'

Figure 322. pSP64 vector promoter and multiple cloning site sequence.
Base #1 is the transcription start site for SP6 RNA polymerase.
The sequence shown corresponds to RNA synthesized by SP6 RNA polymerase.



Use our telephone numbers below.

To order:
800-356-9526
608-274-4330

Technical Services:
800-356-9526



Promega

~~Xho~~ ~~Not~~ ~~Xho~~ ← ~~Xho~~ TK ~~Xba~~

Sylvie had injected w/ 4.9 #13 with XhoI
on SH R37 at concⁿ of 0.1 $\mu\text{g}/\mu\text{l}$
I label this tube B' 35

- ✓ Take 10µl B'35 w/ 49 #13 XhoI (1µg)
✓ + 1µl 10x PBS. B'35

✓ $85-90^\circ$ 5 min

B'35
13 Xhoffteat

✓ +2 p.e (v1 pg) B'32 OS XhoI → XhoI NotI, XhoI

✓ + 2 pl 10x p95 high buffer
✓ + 5 pl 10x p95 high buffer

✓ + 5/2 e d d w

✓ + 1 μ l T4 ligase

15⁰⁰ 328^{pm} to 4:02^{pm}

RT 4.02 ✓ to 4:32 ✓ pm

37° 4:32 to 5:09 pm

2715

B'35
13 Yr old
ART.

Rest to cold man
Fog

Rest
Hofen

✓ 10^6 ($\sim 0.05 \mu g$)

✓ + 80% d.d.w

✓ + 10 μ l 10x PBS

✓ $85^{\circ} \sim 3$ min to ice

B135
#13 Xholing
Diluted

Transformation

1335 DH5 α 1/7

scope

B'35 #13 Xho
diluted

10 pl

5:17^{pm} to 5:46^{pm}

42° 90 sec

Plated

Rent 20

5 20/11

$$5 \overline{) 37}$$

10:50am

Rest
from
here

b

5:57 pm

B34
IK
mini
1-24
37 William
Sent →

(B)



Conclusion The ones with the heavy precipitate were, in general, the poorer ones! But in any case none have become Not I.

Re-inspect oligo & protocol.
Better check ~~post~~ post-ligation
from B'35.

On 12:30pm
 1:42pm 0.86
 MAX HXC4

Mon. Feb 7th

B'35 minus

Processed to KOAc Serum ~ 10am (23 hr.). left until Mon. prior to cent. As usual thereafter - but very low or no ppt. in many tubes. Very odd.

Take 52 μ l (2x26) 10X R1 salts

+ 464 μ l ddw

+ 8 μ l (26x3 units) Not I Lot 714 (0.86)

520 μ l

20 μ l - - - each + 1.5 μ l of minis #1-#24
 37° 10:51 ^{am} to 12:11pm

W

1

2

3

4

5

6

7

8

9

10

11

W

Marker

W

12

13

14

15

16

17

18

19

20

21


22

23

24

W


(3)



This
was
the
trouble

Conclusion:

No obvious reason
for failure - but
gave some times
if not... repeat!
use ϕ 08/H0Q for
Xho most? May need
to prepare fragments
to ↓ ssDNA &
remove dimers & trim



100 μ l of each + 300 μ l EtOH
Back up into 20 μ l 0.001M EDTA

as B38 F_{conc} | B38 M_{conc} | B38 S_{conc}

fragments
made 0.8M
NH₄Ac ~ 85°
1x ϕ 08/H0Q
2x CIAA

Mon. Feb 7th

Xho Not Xho
~~Xho~~ TK Xba

Xho
 Repeat of ligatⁿ → gel check

Take ✓ 36.4 µl ddw
 (≈ 5 µg) → ✓ 7.6 µl w/ 49 #13 super mini (SHR23 0.86 µg/µl)
 ✓ 5 µl 10x R1 salts
 ✓ 1 µl (200) Xho I lot 26
 Now labelled B'39

On 4:40 pm
 of 5:26 pm
 H x 4 0.86
 Max

37° 3.25 pm to 4:30
 37° still

B'39 13 Xho (Post)

1 µl + 16 µl
 B'39 Post

1+16 B'35 w/ 49 #13 Xho I (ex sylvia)
 W
 2+15 B'35 13 Xho ligⁿ (Post)
 W
 Marker
 W
 Dig

Switched

5:55 pm 1x HOQ / HOQ 2x CIAA B'39 13 Xho Post

Take (≈ 2 µg) ✓
 ✓ 20 µl B'39 13 Xho Post
 ✓ + 2 µl 8M NH₄ Ac
 ✓ + 66 µl EtOH

Back up into 14 µl ddw

✓ 14 µl ddw
 ✓ 2 µl 10x p95 ligⁿ buffer
 ✓ 1 µl B3209 X → XNX
 ✓ + 1 µl T4 ligase

B'39
 LIGⁿ

15° 6:46 pm to 6:56
 Cold 6:56 pm to 7:32 am R.T 7:32 to 10:02 37° 10:02 to 10:17

0.8% EtBr

max
On 1:27pm
off 2:38pm

Tues. Feb 8th

Xho Not Xho

Xho

Xba

(could.)

On 7:42pm

0.8% EtBr
max

(1) Dye
~~water~~
F } conc
M }
S } B'38
W
~~water~~

Conclusion: Proceed with M.

Dye	DH52 B'38 (fresh)	80 pl	B'38 M 80 pl	B'38 80 pl
DNA	5 pl F conc	5 pl F conc	5 pl M conc	5 pl S conc

Mini B'38

M #1-24

37° 10:52am to 10:15pm

stationary 37°

Proceeded Thurs. am. as usual

Take 104 ~~pl~~ (26 x 4) pl 10xRS (labeled +1.5M NaCl)
✓ 920 ~~pl~~ pl 10xRS
✓ 16 ~~pl~~ (26 x 60) Not I labeled (100 pl) M & S
To 40 pl, each + 3 pl minus B'38
37° 12:08pm to 1:09pm & back to 37°

B'38 Not 1-24


30/20 pl

Can use excess for Xba I & prep. gel.

5/16 have many
5/16 have some

(B)

(F)



Wed. Feb 9th

• Sol $\boxed{5' \text{ Blank}} \boxed{\text{Eag}}$

Take $\checkmark$ 3 μ l 10 $\times$ R1 salts

$x + 15 \mu l$ of H₂O

$X + 10pe(2pq)$ Paula's 2B2 B'3

$V_1 + 1 \mu\text{e (10V)}$ Eq 1 Lot 22

✓ + $\frac{1}{2}$ p.e (100) Sal 1 lot 47

2pl 1713432

2pe \ 3707 pm


~~Flux 10.10 pm~~

37° $\frac{11.19 \text{ am}}{10}$ to $\frac{12:50 \text{ pm}}{10}$ + 60 pe LdW o to get $\frac{10.10}{10}$

At 1:05 pm

~~3:45 pm~~

Expect
both total
slots



① Conclusion: Forgot that Sal I didn't cut well (see R115 plasmid exp't.). Repeat with 2x more of each enzyme & digest for several hours. Use R115 plasmid.

Or, is it the mg^{-1} ? ~~the mg^{-1}~~

For safety cut out the 9mm-11mm band

~~Dye~~
~~W~~ (2)
~~F~~ ←
~~M~~ ←
~~S~~ ←
~~W~~
~~Marker~~
~~W~~
~~Dye~~

B'43	F
B'43	M
B'43	S

B'43 M



B'43 S

3	M
43	S

018:07pm off 09.55pm
 0.8% Hx42max
 13

0.8% Hx42 max
13

② Conclusion: $\frac{f}{\text{god}}$ Mis $\frac{f}{\text{June}}$!!
 & $\frac{f}{\text{is}}$ $\frac{f}{\text{usable}}$!!






conclusion ① Xba worked fine
judged by NoXba & Xba controls
Fragments prepared




conclusion ② S₀ is terrific

Fri. Feb 11th Ice storm day!

Not Xho ~~~~~ TK Xba



(Fresh) Cell DH5α
DNA



Conclusion ① ligations worked (and there's a DNA's 1). Proportion of TK c'd be 2X and of 5' blank c'd be 4X.

Next time out 370 m up less up time. are diff

Breaked out as B'46 BLUE 1

See also B'59

Processed 5m 2X to 100 pl (Gel B'48)

Plated on 10 plates each 37° 2:30pm Sat

Conclusion

9:15am Sat 1 Blue 1 Procted. Run LG-N gel. Set up a new larger set of transformation & proceed.

(2) Red still zero (2) Blue 4 more


B'46 BLUE 2-5

10:15am Sun 6pm Cold

Processed Mon. Gel B'48 No normal ppts.!

Fri. Feb 11th

Second attempt at NPR1DVP construct



1	Dyp
2	W
3	Pre
4	Post
5	Pre
6	Post
7	W
8	Marker
9	W
10	Dye


B'47 Blue higⁿ

B'47 Red higⁿ

2 pl S
B'45 (NotTKX)

60 pl
B'20 (HindNeoSalM)

75 pl (2 2/3 total)
B'43 M
(Sal5 flank Eag)



504 pl

EtOH

657 pl


Back into 10 pl 1x p95 higⁿ buffer

2+13 pl Pre Blue B'47


2+13 pl Pre Red B'47

2 pl + 13
POST Blue B'47
B'47 Blue
higⁿ

2+13 pl
POST Red B'47
B'47 Red
higⁿ



①


Post
RNAse

conclusion ①: Very odd!
 Whatever it is, it
 has an Xba site

~~and a Sal site~~
 an Xba site

→ several(?) found.


Just repeat ^{digest} with
 more DNA → more times



conclusion ②: Still no
 good. Let's try the
 new ligation → re-
 make 5' flank for
 next try.

Sun. Feb. 13thB'46 Blue 1 test

Take $\checkmark$ 14 μ l (7x2) 10xR1
 $\checkmark$ 115 μ l ddw
 $\checkmark$ 11 μ l B'46 Blue 1
 $\checkmark$ +1.5 μ l (50x6) Xho 1 Lot 26
 $\checkmark$ 5x 20 μ l



37° 11.10am 11.50am

①

Dyp
 W
 No Enz
 Xho
 Xho/Sal
 Xho/Xba
 Xho/Xba/Hind
 Xho/Hind/Sal
 W
 Marker
 W
 Tye

On 12 noon

off 1.0 pm

0.8% Hx4 Max

B'49
digests

Should
 be ~14 kb

On 1.28 pm

off 2.42 pm

0.8% Hx4
Max

10:38am

↓

1:18pm

Dyp

W

1

2

3

4

5

W

Marker

W

Dyp

Dyp

B'46 Blue2-5 test

$\checkmark$ 12 μ l (6x2) 10xR1
 $\checkmark$ 107 μ l ddw
 $\checkmark$ 1.5 μ l (6x50) Xho


5x20 μ leach μ l10 μ l mini

2-5

2 μ l mini 1B'49²

1-5

Xho



① Conclusion:
Same ligated
products are
visible.

Proceed to
prepare more
5' flank


① Cyn 4:10pm
off 4:40pm

1	Blue
2	Red
3	W
4	Marker
5	W
6	Dye

10:53am B51 Minis 1-12 BLUE ← Picked
13-24 RED ← easily
Picked only
with difficulty.

6pm to stationary

Mini #24 re-seeded from plate
at 6:35pm to shake - it was not
growing. #21 is zero on both.




Hx4
0.8%Sun - Feb 13th3rd (New) titration for NPR1 Duphit
hit } B'51

Take

✓ 1 μ l B'45S Not TK Xba
 ✓ 30 μ l B'20M Hunk Neo Sel

~35 μ l B'43M (Rest) Sel 5' flank Eaq



Cold 6.10 pm

Sunday

3:25 pm Monday

Blue

Red

Marker

Fresh H 5x 1/7 150 μ l 150 μ lBlue hq 1 μ l 1 μ l Red hq0 $^{\circ}$ C 3:25 pm to 3:55 pm90 $^{\circ}$ C 42 $^{\circ}$

Pre 1 B'55 Blue

Post 1 hq

Pre 2 B'55 Red

Post 2 hq

W Dye

32 $^{\circ}$

4:13 pm Pooled

5/16

20/16

5, 20, 40, 40, 40

5




F
M
S



Conclusion: Mufine - processed.
Total amount available
is roughly 10X that
variable in gel.

Mon. Feb 14th NM day
 Re-make of Sal 5' Flank Eag



On 6:28 pm

off ~ 10:30 am

1 v/cm 100 mA


B'43'
2B2 En/Sal

↑ to 125 mA at 9:45 pm

Processed to → B'53 F
 Used whole → B'53 M
 Discarded → B'53 S

	Dye	W
Undiluted	F	B'53
	M	
	S	
On 1 pm	W	
Off 1:45 pm	Marker	
Max	W	
0.8% Hx4	Dye	

	B'54 BLUE	B'54 RED
DH5d 1/7	80 μ l	80 μ l
B'55 LIG ^W DNA	2 μ l	2 μ l
	0° 5:40	to 6:10 ✓
	42°	90 sec ✓
	Plated	NZY 1 Amp
	Rest 20, 5, 20, 5	16, 16
	37° 6:25 pm	



↑
B'51 gel

Conclusion:

Band proportion was quite good. Cancels the ligated products. Go ahead & repeat transformation with B'55 BLUE & RED ligations - ~ 3 μ l are left

Meanwhile streaked out the B'54 BLUE #1

Also have another look at B'46 BLUE 1 & possibly pick more of B'51.

Run gel & trouble

See B'51

See B'64 3
for B'55 Red

See B'57
-renamed

B'54
Blue Rest

Only 1 on ~~RED~~ Blue Rest
0 on ~~Red~~

B'55³
RED

80 pl

Rest
≤ 2 pl

11:50am

Tue. Feb 20

~ 12:30

2.5

5


10x7

12:50pm

Tues

↓
Processed
ThB'55
A131Take
B'53
MTues Feb 15th

Third attempt at NPR1 DVP construct

B'55 Blue
highB'55 Red
high

- ✓ 8 pl S B'45 Not (TK) Xba
- ✓ 60 pl M B'20 Hind Neo Sal
- ✓ Whole M B'53 Sal (5' flank) Eco

~130 pl ~130 pl

✓ 100 pl B'11/148
Blue150 pl B'11/155
RedSame
baggiesExtracted 1x H₂O/H₂O

190 pl / 230 2x CIPA
+ 570 ~~1000~~ pl. EtOH Transferred to new tube ~250 pl / 290
750 pl EtOH

Back into 10 p p 95 1x high buffer 210

✓ 2+13
B'55
P1C Blue

✓ 1/2 1/16/16
Lot 11005 1/2 pl + 1/2 pl
TK ligase TK ligase

✓ 2+13
Red B'55
P1C

✓ 2+13
B'55
Post. Blue

15:0 4:37
RT 5:08 to 5:23
37° 5:23 to 5:38

✓ 2+13
Cold Red B'55
Post



Conclusion: B'51² BLUE #32 (see B'61)
 looks v.v. promising.

B'51² BLUE

see B'61)

On 8:46am
off 9:46am
0.8/10 HFE MAX

Wed. Feb 16th

Re-transformation of B'55 ligation

1	
2	
3	
4	B'57 BLUE
5	
6	
7	
8	
9	
21	B'57 RED
23	
32	B'51 ² Blue
W	
Marker	
W	
Byte	

	B'55 ² BLUE	B'51 ² BLUE	B'55 ² RED	B'51 ² RED
DH5α	125 µl	125 µl	125 µl	125 µl
DNA LIGase	3 µl	3 µl (REST)	3 µl	3 µl (REST)
0°C	3:08 pm	3:21 pm	3:44 pm	3:05 pm
2:33 pm				
	42°	90 sec		
Plated	1.5 µl	6 µl	250 µl	4
	37°	3:35 pm		



Re-names:

B'54 BLUE	
1	≡ B'54 BLUE #1
2	≡ B'51 ²
3	≡ B'51 ²
4	≡ B'51 ²
5	≡ B'51 ²
6	≡ B'55 ²
7	≡ B'55 ²
8	≡ B'55 ²
9	≡ B'55 ²

B'57
Minis 1-9, 21-23

37° ~ 10:30am ~ 7:30am

#22 Red did
not grow



Conclusion:

The old result
is not changed
appreciably
this is worse!

unrelated 1359

Wed. Feb 16 d

B'46 Blue 1 tests (repeated)

On 1:53 am
0.80 1x4
NAT

Water

NAT

Xho

X/Sal

X/Xba

X/Xba/Hind

X/Hind/Sal

Marker

W

Eye

✓ Take 14 pe (7 x 2) 10 x R 1

✓ 112 104 pe 1 w

* 14 22 pe B'46 Blue 1

+ 1.5 pe (50 x 6) Xho 1 lot 26

20 pe

B'59/1
No Enz

1 pe ✓
Sal 47

1/2 pe ✓
Xba 49

1/2 pe ✓
Xba 45

1/2 pe ✓
Hind 45

1/2 pe ✓
Sal

All are
20 pe

B'59/1
Xho

B'59/1
X/Sal

B'59/1
X/Xba

B'59/1
X/Xba/Hind

B'59/1
X/Hind/Sal

37° 11:53 am — 1:40 pm
Thurs.


* Can't find the mini. One (v. large) colony ∴ into 50 ml 1x AMP NZY overnight 5:50 pm

B'46²
BLUE 1

Back into 100 pe cells etc.

Looked good & it is single colony

+ changed to 14 pe B'46² BLUE 1



Conclusion: At least BLUE


Red safe 1 with
B'512 #13, 16, 23, 30 & 32

all large & the same.

11:30am Processed as usual to 50 μ p

FR

B'512
BLUE
13, 16, 23, 30, 32
(contd.)



On 9:50am
 off 11:10am
 0.80 HX4
 750mt

Thurs. Feb 17th

Rescue of last available colonies B'51

The Blue & Red plates from B'51 are still available* with very solidified colonies - often with no visible parent colony.

Following were rescued on 2xAMP N2Y plate

2xAMP cultures

Not all
 are single
 colonies

B'51²
 13
 14
 15 (Red)
 16
 17

B'51²
 13 -
 -32 (Blue)

37° 4:50pm
 Th.


B'51²
 BLUE

32th on B'57 gel

* from the garbage
 pail!

Processed to KOAc on Fri. Feb 18th
 estimated 9am.

W
 Marker
 W
 Dye



Conclusion: These look v. good - but there must be a 1.1 Kb band under the RNA in X/Ting/Sel lane. Checking after RNase confirms OK.

Streaked out v.6 pm


B'51² 30

as

NPR1 DUP
BLUE B'62

Removed
BLUE 3'
B'62
see B'73

Orientation needs checking with PCR versus the 3' flank phage. Make RED only if orientation checks correctly.



→ Poor tiny colonies.

Re-streaked out on fresh NZY 1X AMP plate

also B'51²

13, 16, 23, 32
30

5 pm
Sat

All colonies are small, but set up 100 ml of one colony from NZY

1X Amp

B'51²

30


10:45 am Sun

→ To rt. 11:40 am Sun


Fre. Feb 18th

NPR1 DVP Blue tests


Take ✓ 24 μ l (12x2) 10xR1
 ✓ 213 μ l ddw
 (12x5) ✓ 3 μ l Xho 1 Lot 26



13 } B'63
 16 } Xho digests
 23 } undig
 32 }
 30 }
 30 X/Sal B'63
 30 X/Xba digests
 30 X/Xba/Hind undig
 30 X/Hind/Sal
 W
 marker
 W
 dye



Processed to KPAc - all had ~ normal (low) ^{but}
looking pellets.



Conclusion: None are correct!
Try rest of B'55 (as B'55³)
and 2 pl more of B'65
(as B'65²)


Did so. 3 colonies B'65²
from B'65² [217, 218, 219]
1 colony from B'55³

[RED] B'55³
31

2X Amp
To 37° as mixed ~ 3:20pm

Tues. Feb 23rd
7 streaked out
on NZY 2X Amp

~ 9:30am Th highly
growth but all OK



Conclusion: looks as if B'55³ #31
is good, see B'65.

(2) Dye On 10:20
W Off 10:50
217 } B'65²
218 }
219 }
31 B'55³
W
Marker
W
Dye

B'65 mixing

Sat. Feb 19th

Repeat of NPR 1 DUP RED higⁿ
(B'51)

10:02am
11:04am
0.56 HX4 MAX

- ✓ Take 1 µl B'45 S Not TK Xba
- ✓ 30 µl B'20 M fluid Neo Sal
- ✓ Whole (w/150 µl) B'43 F
- ✓ 75 µl B'11 155 Red 3' flank

B'43
F

+ 768 µl EtOH

Back into 10 µl 1x p95 higⁿ buffer
+ 1 µl 1 µl + 4 µl p95 JL tube
Th ligase

Possibly
a little
more

Cold 6:20 pm

B'65
RED
HIGⁿ

10:50am Sun

B'652	B'65
✓ 80 µl	✓ 200 µl
✓ 2 µl	✓ 4 µl
DN5d 1/7	B'65 Red hig ⁿ
Tues 11:56am	0° 10:53am
Feb 22 12:30	11:24am
90sec	4.2°
plate of N2Y 1X AMP	
Total (w/12)	
2, 4, 8, 16, 20, 20, 20, 20 Rest	

B'65
2

B'65
Rest

2.5, 5, 10 x 7
12:50 pm Tues

37° 11:40am Sun

12 colonies (all of variety but #4 & #15) to minis

9:24am Mon → 5pm → 9am Tues

9am Mon

5 µl to new replica plate

20
11:50
0.56 HX4
25
53
Sun
10x
2x

Wed Feb 22nd

Using up last of B'65 high

DH5
1/7

DH5x 1/7	240 pl	DH5x 125 pl 10.3 025
high	B'65 Red high ~4 pl	3 pl B'65 Red high
0°	4:04 pm	4:34 pm
42°	4:45	90 sec
Plated		1.25, 1.5, 2.5, 10, 20 -
(25) 20 - - - -		(13)
	37° ~ 5 pm	

DH5
10.3

B'65
RED HIGH

N 9:30 am


Stored in cold
- pending analysis

B'55 Red 31

Thurs. Feb 24th

Discarded


Streaked out
for Sylvia on
suitable plate
Minimal medium



Conclusion:
Self-evident

On 3:59 pm
0.85.10
0.85.10

Bye
W
Uncut
Xho
X/Sal
X/Xba
X/Xba
X/Hind
W
Marker
W
Bye



5:59 pm
 5:10 pm
 0.89 HXB

max



Thurs. Feb. 24th

Red

69

NPR1 DUP Red B'55

B'p
 W
 Uncut
 Xho
 X/Sal
 X/Xba
 X/Xba
 X/Hind
 W
 Marker
 W
 Dye



B'69
 Xho

X/Sal

X/Xba


X/Xba
 Hind

X/Hind
 Sal

37° 12.45 pm → 3:45 pm

11

11




Conclusion:

The Pst patterns are identical (see * below) but the Pst/Xba differ & allow orientation to be determined, but beware of PCR using any promoter primers

SHS 154
B'553 Red
31
1.275 µg/µl

**RED
31**

5' flank




* Same site because both use PstI promoter & enhancer

600 → 2x300 with Xba
1300 does not change

**BLUE
30**

5' flank



1300 → 300 + 1000 with Xba

600 does not change

SHS 148/158
Blue Dup NPR1
B'62
0.775 µg/µl

Rename B'62 Blue 30 see B'73

on 11:42
off 12:47
0.86 HX4 MAX


RED
31

BLUE
30


Fri. Feb 25th

Cross-checks of Red v. Blue

Age
W
Pst Blue
Pst Red
Pst/Xba Blue
Pst/Xba Red
W
Marker
W
Age




2 poss. orientations
- other Pst sites unknown



37° 9:54pm
11.29am


See PMO 3-127.
The desired orientation for duplication is RED 31

①



①
Conclusion: B'62 BLUE 30 has
 Xho as expected, but too many
 Eco sites, including one close
 to Xho. Try Not I (w/ 1 in 4
 chance adding none elsewhere)

②



②

Conclusion: B'62 BLUE 30 has
 the Sal site expected & generally
 the correct Sal/Xho fragments.
 But there is no Not I site.
 Can proceed by changing
 Xho to Xho Not Xho again
 & reinserting Sal Eco?
 { Reversed }
 into Sal Not.

Sat. Feb 26th

BLUE
30

Conversion of DUP to DEL(etc)

Can extract the sal $\xrightarrow{5 \text{ Blank}}$ Eag fragt

from the incorrect Red or Blue & invert it by site switching - a many other possibilities. Check if Eag sites unique (still present).

Sylvia prep of B'62 BLUE NPR1 DUP - now

B'62 Blue 30

11:55am
12:28pm
0.8pg
Hx4 Max

①

Agg
W
Uncut
X
XE
E
W
Marker
W
Dye

Take $\checkmark$ 8 pl (4x2 pl) 10x R1 salts
 $\checkmark$ 71 pl ddw
 $\checkmark$ (0.8pg) 1 pl B'62 BLUE 30

Rest $\checkmark$ 20 pl $\checkmark$ 20 pl $\checkmark$ 20 pl

$\frac{1}{2}$ pl Xho I Lot 26 $\frac{1}{2}$ pl Xho $\frac{1}{2}$ pl Eag I Lot 22

Uncut B30 $\checkmark$ B30 $\checkmark$ XE B30 $\checkmark$ E B30

37° 10:43am $\rightarrow$ 11:45am

②

Agg
W
Not (N)
Xho (X)
Not + Xho (NX)
W
M
W
Dye

5:20pm
6:06pm
0.8pg
Hx4 Max

Take $\checkmark$ 10 pl (5x2 pl) 10x R1 salts
 $\checkmark$ 89 pl ddw
 $\checkmark$ 1.2 pl B'62 BLUE 30

Rest $\checkmark$ 50 pl $\checkmark$ 10 pl $\checkmark$ 10 pl $\checkmark$ 20 pl

$\frac{1}{2}$ pl Not I 46 $\frac{1}{2}$ pl Xho 26 $\frac{1}{2}$ pl Sel I 26

N B30 $\checkmark$ NS B30 $\checkmark$ X B30 $\checkmark$ XS B30

3:40pm $\rightarrow$ 5:10pm

Check on Not I site

Forward

Ace Summary

ACE 1 (rev +): 6/11/93

DNA: SHR25 w110 #7 xhoI
2 Electroporations 3ul DNA and 30ul DNA
Pick 40 colonies and froze 20 colonies
2 vials per colony.

Positive control
for ACE-DVP

ACE 2 (rev +): 6/25/93

DNA: SHR35 w157 "8"
2 Electroporations 3ul DNA and 30ul DNA
Pick 20 colonies. Froze 2 vials (or more)
from 16 colonies.

Positive control
for ACE-DVP
with Neo reversed

This is
the correct
one to use.

ACEDUP-1: 7/10/93

DNA: SHR59 Rev6, A'19 Rev 10ug SH
1 Electroporation with 10ug DNA
Pick 180 colonies, all negative.

ACE-DVP
with Neo reversed

ACEDUP-2: 7/14/93

DNA: SHR59 Rev6, A'19 Rev 10ug SH
1 Electroporation with 10ug DNA
Pick 120 colonies, 8 positive lines.

ACE-DVP
with Neo reversed

Cell line	# vials frozen	chromosome % at 40	injection date(s)	result
-----------	----------------	--------------------	-------------------	--------

7C	12			
15A	8	0%, 70%		
6D	10	80%		
2B ← S	8	62%, 100%		

→ 8/23/93 2 chimera - weak
8/25/93 no chimera
9/8/93 Not pres.
9/9/93 No chimera

← only
and
from
here

3B ← S	11	66%, 80%		
3D ← S	7		10/6/93	Not pres
5A	12	80%		
4B	8			

A
if
external glow.
mean
turn OK

Thaw 3D for chromosome

Thaw 6D for Southern

But it is really
 $AGT \rightarrow ANG I$
not total
removal

which must
equal production

suspicious
loop

Th

Tues. March 29th

Some numbers on renin / angiotensin

Assumptions:

- (i) 1, 2, 3 & 4 gene copy animals are to a first approximation normotensive.
- (ii) all corrections are via renin conc. changes.
- (iii) Angiotensin II production is constant (corollary of assumption (i) & (ii))
- (iv) Angiotensin II production changes linearly with angiotensinogen and/or renin conc.^{ns}
- (v) the duplication chromosome is not twice normal.

Normal $2p - u = 100$

1/0 $p - u = 35$

$\therefore p = 65$ and $u = 30$

(p = production from one normal chromosome.
 u = utilization of AGT)

2/1 $x \cdot 2p + p - u = 124$

$\therefore x = 0.685$

(x = fractional efficiency of an AGT gene in duplicated case)

Predicted 2/2 $2 \cdot x \cdot 2p - u = 148$

Predicted 2/0 $x \cdot 2p - u = 59$

This looks good $\rightarrow$ $\begin{bmatrix} 100 \end{bmatrix} \times R_2 = 100\%$ $R_2 = 1$ (assumed)
 $\left\{ \begin{array}{l} \text{Normal AGT level} \\ \text{Normal Renin level} \end{array} \right\}$ ATI production

$\therefore$ Predicted 1/0

$R_1 = 2.86$

Predicted 2/1

$R_3 = 0.81$

Predicted 2/2

$R_4 = 0.68$

Predicted 2/0

$R_{2/0} = 1.69$

Predicted 0/0

$R_0 = \text{off scale}$

If normotensive

Wed. April 6th

Another try!

let clearance be prop^l to [AGT]

let $u_{100} = \dots$ [AGT] constant for 1, 2, 3

let 2AGT clearance be 2x normal

$$1/1 \quad 2p = u_{100} + c_{100}$$

$$1/0 \quad p = 0.35 u_{100} + 0.35 c_{100} = 0.35 (u_{100} + c_{100})$$

$$2/1 \quad 2px + p = 1.24 u_{100} + 1.24 c_{100}$$

2) are inconsistent.

Try u constant (for normal sensitivity)
with clearance prop^l to [AGT]

$$1/1 \quad 2p = u_{100} + c_{100}$$

$$1/0 \quad p = u_{100} + 0.35 c_{100}$$

$$2/1 \quad 2px + p = u_{100} + 1.24 c_{100}$$

$$\left. \begin{array}{l} 1-2) \\ p = 0.65 c_{100} \end{array} \right\}$$

$$\text{and } u_{100} = (0.65 - 0.35) c_{100} = 0.3 c_{100}$$

$$\text{or } c_{100} = 1.54 p \quad \text{c.f. B'79}$$

$$3) - 2) \quad 2px = 0.89 c_{100} = 0.89 \times 1.54 p$$

$$\therefore \bar{x} = \frac{0.89 \times 1.54}{2} = 0.6853 \quad \text{c.f. B'77}$$

trial:

$$2/0 \quad 2px = u_{100} + q \cdot c_{100} \quad \leftarrow \text{AGT level}$$

$$\therefore 0.89 c_{100} = 0.3 c_{100} + q \cdot c_{100}$$

$$\text{and } q = 0.89 - 0.3 = 0.59$$

AGT 2/0

59%

Remain levels are

$$1/1 = 1.0$$

$$1/0 = 2.86$$

$$2/1 = 0.81$$

$$2/0 = 1.69$$

Predictions

Sat. April 18th

Further numbers

Assume production is just linear with gene dosage;
 Assume 1, 2, 3 copy animals are nonmotensive &
 "therefore" make same amounts of AT I
 Assume non-specific clearance is proportional to [AGT].

1) $1/1$ $2p - \text{clearance at 100\% plasma level } C_{100} = \text{utilization to make AT I} = u$

2) $2/1$ $3p - 1.24 C_{100} = u$

3) $1/0$ $p - 0.35 C_{100} = u$

From 1) - 3) $p = 0.65 C_{100}$ or $C_{100} = 1.54 p$

But from 2) - 1) $p = 0.24 C_{100}$

This is not possible.

All of which makes the clearance (??) receptor of ANF more interesting.

2/2 5) $4px = u_{100} + r C_{100}$

But $p = 0.65 C_{100} \rightarrow u_{100} = 0.3 C_{100} \rightarrow x = 0.6853$

$\therefore r = 4 \times 0.65 \times 0.6853 - 0.3 = 148\%$ 2/2

AGT

With renin

$2/2 = 0.68$

600153
4/26/94PRODUCT: **Cloned *Pfu* DNA Polymerase**

CATALOG #: 600153 600154 600159

QUANTITY: 40 μ l 100 U 500 U 1000 UCONCENTRATION: 2500 U/ml 250/ μ l

SPECIFIC ACTIVITY: 75,000 U/mg

LOT #: **163006F**CERTIFIED BY: Mila Angert *M. Angert*

DATE: April 11, 1994

QUALITY CONTROLLED BY: Jeff Amberg *Jeff Amberg*

DATE: April 14, 1994

DESCRIPTION: Cloned *Pfu* DNA polymerase is isolated from the hyperthermophilic marine archaeobacterium, *Pyrococcus furiosus*. The multifunctional thermostable enzyme possesses both 5' to 3' DNA polymerase and 3' to 5' exonuclease activities. The 3' to 5' proofreading activity associated with cloned *Pfu* DNA polymerase results in a 12-fold increase in fidelity of DNA synthesis over *Taq* DNA polymerase. Cloned *Pfu* DNA polymerase has a temperature optimum between 72–78°C, and remains greater than 95% active following one hour incubation at 95°C.

REACTION BUFFERS: **10x Cloned *Pfu* Polymerase Reaction Buffer***
200 mM Tris-HCl (pH 8.75)
100 mM KCl
100 mM (NH₄)₂SO₄
20 mM MgSO₄
1% Triton® X-100
1 mg/ml BSA

REACTION CONDITIONS: **Note: All reagents, including cloned *Pfu* DNA polymerase, should be mixed immediately before use.**

DNA synthesis is performed in 100 μ l of a mixture containing 20–200 μ M dNTPs, 0.3–1 μ M primers, 0.1–250 ng template DNA, 10 μ l 10x polymerase buffer and 2.5 U *Pfu* DNA polymerase. Mix the reaction gently, centrifuge briefly, then overlay with light mineral oil. Initially, denature the reaction by incubating at 95°C for five minutes; then cool to 40–68°C for 5 minutes to allow primers to anneal to the template DNA; then incubate at 72–78°C to initiate primer extension.

*Formerly sold as Buffer #3.

True.

Tues. April 26th

Simple test of Wayne Barnes' principle with ATR1B amplification

Conditions for 1.2 kb PCR have been border line
 $\therefore$ ~~best~~ $\pm$ editing Tag (Pfu) - using $\frac{1}{10}$ the vol. (intended 1) Barnes 20

Usual formula (see A'121):

Today n=12	ddw.	$n \times 2.85 \mu l$	
34.2 ✓	pH 8.8	20x PCR A'121	$n \times 1.25 \mu l \equiv 67 \text{ mM}$
15 ✓		4 dNTP's 25 mM	$n \times 0.1 \mu l \equiv 250 \mu \text{M}$
1.2 ✓		DMSO A'121	$n \times 2.5 \mu l \equiv 10\%$
30 ✓		Kim's w 139 3' Neoprimers	$n \times 0.3 \mu l$
3.6 ✓		A'102 ATR1B primer	$n \times 1 \mu l$
12 ✓		Standard Tag A'121/	$n \times 2 \mu l$
24 ✓		+ B'81	$n \times 10 \mu l$
120 μl			

This is 18 Tag
 10 by volume
 Rest 1.4 μl Pfu B'81
 + $n \times 15 \mu l$ DNA sample
 16 mM $(\text{NH}_4)_2\text{SO}_4$
 6.7 mM Mg^{2+}
 16 mM $(\text{NH}_4)_2\text{SO}_4$
 3.5 mM Mg^{2+}

PCR mix (modified from the recommendations of New England Biolabs) is 67 mM Tris-HCl (pH 8.8), 6.7 mM MgCl_2 , 16.6 mM $(\text{NH}_4)_2\text{SO}_4$, 10 mM 2-mercapto-ethanol, 6.7 μM Na_2EDTA , 10% dimethylsulfoxide, 330 μM of each dNTP, 3.0 μg of each primer, 1 unit Tag polymerase per 50 μl .

$\therefore$ modify by $\pm$ $n \times 0.1 \mu l$ of Pfu

Positive controls are { Pos. Control A109 6E and A109 12D } ~~use 2 μl~~

Finished $\rightarrow$


try 5 μl (A11-4 7B)
 Try A11-4 16C - these are real clones
 Use 15 μl

Back	Temp	Time	
35 cycles	95°	92°	Total 1 min
	39°	60°	Total 7 sec
	60°	60°	10 min

A11-4 16C - this is the +ve pool
 use 15 μl
 On 5.15 pm 1st cycle not v. good
 are temps.

6E
 2D
 7B
 16C
 16
 6E+
 1D+
 7B+
 6C+
 16+

Today	Back	Temp	Time	
1	10 μ l	5 μ l 6E	2 μ l 12D	✓
2	13 μ l	2 μ l 12D		✓
3	10 μ l	5 μ l 7B		✓
4	10 μ l	5 μ l 16C		✓
5	5 μ l	10 μ l 16		✓
6	10 μ l	5 μ l 6E	2 μ l 12D	✓
7	13 μ l	2 μ l 12D		✓
8	10 μ l	5 μ l 7B		✓
9	10 μ l	5 μ l 16C		✓
10	5 μ l	10 μ l 16		✓



On 10:50am 10V/cm 300mA 2% Ag Hx1
Off 11:28am

83

Dye		
Water		
1	-	5pl A109 (6E)
6	+	
2	-	2pl A109 (12D)
7	+	
3	-	~5pl A'III-4 (1B)
8	+	
4	-	5pl A'III-4 (16C)
9	+	
5	-	10pl A'III-4 (Pool)
10	+	
W		
Ag Marker		
W		
Dye		

Conclusion: The Pfu definitely can be beneficial. However 7B & 16C show the unexplained failure mode. Simon thinks that too many cycles can cause this. Wayne says it occurs more frequently when larger amounts of enzyme are being used. ∴ test both variables - less enzyme & fewer cycles. Don't have much sample left ∴ develop technique with ~~the~~ ES cell DNA isolated by Damić from our current infection cell lines.



No Pfu

1 μ l DNA
(~50ng)

~100ng

~150ng

16C

M9:38am
of Yellow by to
wick

~1.6C

+ Pfu

~100ng

16C

~150ng

Conclusion: Clearly with control the +Pfu is better
but linearity with DNA conc. is lost

At 111-4 16C is positive in two tests
three

Clear that the problem
is likely to be linked
by $\downarrow$ DNA & the Pfu helps.
Pfu does no clear help.

Try again with 5ng and 10ng of DNA + Pfu
Also try with NaOH denat.

Control 1 μ l
2 μ l
3 μ l

Control 1 μ l
2 μ l + Pfu
3 μ l

1
2
3
4
5
6
7
8
9
10
11
12
1'
2'
3'
4'
5'
6'
7'
8'
9'
10'
11'
12'
13
14
15
13'
14'
15'

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

1 μ l
3 μ l

to
week

~1.2% ~ HX2
600mA

87


Thurs April 28th

More PCR tests

A111-4 16C and A111-4 B
1.1 $\mu\text{g}/\mu\text{l}$ 1.06 $\mu\text{g}/\mu\text{l}$




Take 200 μl of each. 1x HOG / HOG
2x CIAA

Make: —



~~16C~~ 16C (phenol) at 0.05 $\mu\text{g}/\mu\text{l}$ etc

i.e. 195 μl ~~COOH~~ T/E buffer
+ 3.1 μl ~~16C~~ or 3.1 μl ~~16B~~ + 4.7 μl 4B or 4BP



Set up for 30 samples

3:38 pm	Start
95 92.5°	1 min
40 ~61°	7 sec
60 60°	10 min
Good load at 60°	

ddw 536 μl ✓
20x PCR 37.5 μl ✓
4 dNTPs 3 μl ✓
DMSO 75 μl ✓
W1393 Neo primer 9 μl ✓
A102 ATRIS primer 30 μl ✓

$$30 \times 23 = 690 \mu\text{l}$$

+ 12 μl Tag ✓
Rest ~~12~~ μl → 345 μl + 0.6 μl Pfu B'81

0.40 / 25 μl

29 μl - Pfu								29 μl + Pfu							
1	2	3	4	5	6	7	8	1'	2'	3'	4'	5'	6'	7'	8'
1 μl 16C	16CP	16B	16BP	16C	16CP	16B	16BP	ditto							
9	10	11	12	13	14	15		9'	10'	11'	12'	13'			
3 μl 16C	16CP	16B	16BP	16C	16CP	16B	16BP	16C	16CP	16B	16BP	16C	16CP	16B	16BP



2:12:07 pm
~~2:12:07 pm~~

	W	
2' pl	1	C/2
	2	C/4
	3	C/8
PSU	4	B/2
	5	B/4
	6	B/8
	1'	C/2
	2'	C/4
PSU	3'	C/8
	4'	B/2
	5'	B/4
	6'	B/8
Note reverse order of above	30	Alkali B
	31	denat C
	30'	B
	31'	C
	W	Marker

Conclusion: Went backwards!
 Return to 1st pl of Tag

For alkali start
 should really hold
 60° for 10 min.

5:20 pm

Good 60° followed
 by 92° start
 max temp

1:00 93°

7 sec ~ 61°

10:00 60.7°

35 cycles

Start hold 10 min.	B' 87	Bye
✓	16 C	7 B
✓	50 mg/pl	50 mg/pl
✓	10 pl	10 pl
✓	10 pl TE	+ 10 pl TE
✓	B' 88	B/2
✓	C/2	
✓	5 pl	5 pl
✓	15 pl TE	15 pl TE
✓	B' 88	B/4
✓	C/4	
✓	2.5 pl	2.5 pl
✓	+ 17.5 pl TE	+ 17.5 pl TE
✓	B' 88	C/8
✓	C/8	
		2 pl
		2 pl

1.4% HX 1 ~ 8V/cm

Tri April 29th

PCR Contd.

May need revision to keep salt conc. lower this will give n final

Test standard method $\pm$ Pfu with DNA levels decreased still further. Also try alkali denat. for step 1 :-

Take 4 μ l of B'87 ^{50ng/ μ l}
+ 2.6 μ l of { 8 μ l A'102 ATR1B primer
+ 0.5 μ l W139 3'Neo primer

7B ✓	16C ✓
4 μ l	4 μ l
2.6 μ l	2.6 μ l
✓	✓
3.3 μ l	3.3 μ l
✓	✓
3.3 μ l	3.3 μ l
Sat at room temp. for several hours	Sat at room temp. for several hours
B'89B (3.2 μ l)	B'89C (3.2 μ l)

B'89 ALKALI

300 μ l 2M NaOH

700 μ l dH₂O

B'89 ACID

300 μ l 2M HCl

200 μ l 1M Tris pH 8

500 μ l ddw

+ 3.3 μ l 0.6M NaOH $\rightarrow$ pH 12.5 minutes

+ 3.3 μ l 0.6M HCl $\rightarrow$ pH 8

+ 0.2M Tris HCl pH 8

Use 8 μ l in $\pm$ Pfu with Special Mix n=5

Regular Mix n=14 ddw 250 μ l (n x 18 μ l)
20 x PCR 17.5 μ l (n x 1.25 μ l)
4 dNTPs 1.4 μ l (n x 0.1 μ l)
DMSO 35 μ l (n x 2.5 μ l)
W139 3'Neo primer 4.2 μ l (n x 0.3 μ l)
A'102 ATR1B primer 1.4 μ l (n x 1 μ l)
0.4 μ l / 25 μ l $\rightarrow$ 32.2 μ l (14 x 2.3 μ l)
+ 5.6 μ l Tag 1

ddw 64 μ l
20 x PCR 6.25 μ l
4 dNTPs 0.5 μ l
DMSO 12.5 μ l
Standard Tag 2 μ l (0.4 μ l / 25 μ l)
(5 x 17 μ l) 8.5 μ l

22

23 μ l

rest

100 μ l
+ 2.3 μ l Pfu

B'89B (30) B'89C (31) B'89B (30) B'89C (31)
n 8 μ l n 5 μ l n 7 μ l n 6 μ l

Unequal ✓

1 2 3 4 5 6
C C C B B/2 B/4 B/8
2' 3' 4' 5' 6' 12.5 μ g

EXPERIMENT #1

AGT TAILCUFF DATA

Sheet 12

COMPILATION

SUMMARY OF ANTHOLOGY DATA

10 COPY ANIMALS														
MOUSE	8667	8739												
EXP	1	3												
SEX	F	M												
BP	101	91												
HR	736	790												
1 COPY ANIMALS														
MOUSE	8666	8668	8678	8679	8680	8689	8687	8688	8725	8726	8740	8741	8771	
EXP	1	1	2	2	2	2	2	2	3	3	3	3	3	
SEX	F	F	F	F	F	F	F	F	M	M	M	M	M	
BP	125	138	106	96	100	108	108	105	105	119	112	104	116	
HR	700	750	705	689	707	663	745	784	788	737	777	791	748	
MEAN 1	131.5	725												
STDEV 1	6.5	25												
SE 1	4.596194	17.67767												
2 COPY ANIMALS														
MOUSE	9204	9206	9248	9256	8690	8738	8753	8754	8776	8777				
EXP	1	1	2	2	2	3	3	3	3	3				
SEX	F	F	F	F	F	M	M	M	M	M				
BP	129	123	132	99	82	125	133	112	120	126				
HR	701	657	683	712	740	716	815	743	774	769				
MEAN 1	126	679												
STDEV 1	3	22												
SE 1	2.12132	15.55635												
3 OR 4 COPY ANIMALS														
MOUSE	8657	9247	8646	9253	9200	9250	9199	8647	9254	9249	9255			
EXP	1	2	2	2	2	2	2	2	2	2	2			
SEX	F	F	F	F	F	F	F	F	F	F	F			
BP	171	118	97	106	124	124	109.3	134	122	131	140			
HR	704	716	663	730	690	704	711	687	740	700	707			
MEAN 2														
STDEV 2														
SE 2														

RANK #1

#2


#3

#4

May 9th, 1994

9 references

1205 200.1566
704.8 etc



5'

note: D8Mit56 Left

3'

A	C	A	C	T	C	A	G	A	G	A	C	C	A	T
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

24
50%

G	A	G	T	A	C	A	C	C						
---	---	---	---	---	---	---	---	---	--	--	--	--	--	--

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

5'

note: D8Mit56 Right

3'

G	A	G	T	T	C	A	C	T	A	C	C	C	A	C
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

L34 D8Mit14

Tue 27 Apr 1993

MAP POSITION: 64.9 cM from top of Chr8 (Framework)

PRIMERS: TTTTCACACTCACGTGTGCG

GTCTCTCCTTCCTGGCGCTG

PRODUCT: 143 bp

GENOTYPE: HBABBAHHAHBAHHHHHHBA-AB BAHABBABBBHHHAHBHHHHHH (rank 1)

SEQUENCE:	10	20	30	40	50	60
0	GGGTGCTGGT	GGAGNCCTCG	NATACCCGGG	NGATCTAAGA	GGCTCTTCTA	GACTTTGTAG
60	GTACCTGTTT	TCACACTCAC	GTGTGCGTAC	AGACTGAAAC	ACACACACAC	ACACACACAC
120	ACACACACAA	TTAGAAATAA	TACTGAAAAA	CAGAACAGGA	AGTTGGCTGA	AGTATACGTG
180	CGTGTTACCC	CAGCGCCAGG	AAGGAGAGAC	GGTGGATCAT	CCATGCACAC	GGCCCCATGA
240	TAAAATATCA	NGCTTATGGA	ANGNGTTCAG	TTTNTCTAAA	CCACTGGGNN	NATTTTCAGTC
300	TATGTGCTTG	TGTCTGTANT	AGAAGGNNGT	NANTATCAAC	TAACTAAAGA	TTTCTTGTTG
360	TTTAAAAAAG	GNGGNANGGN	GNCCTGGGGT	CAGAACTGCA	TGTACAANAG	CCCATTNTCT
420	CCAGNGACTN	TCAANCCCCA	AGNTGCCCTA	GCATNGCCAN	NGTTTTGAGG	ATTCCTCTNA
480	GAGTCCGCCC	TTNNGNCNTT	NANGGTTTGG	CGTANTNGTG	NGTGCATANT	TTTTTCCTGT
540	GTAAATTGT	TNTNNGTNNN	ANTTTNCACA	NNCNTNNGG	GCCCGAGAGT	TNNNGTNTGG
600	NGC					

ALLELES: ob cast spr a b6 c3h dba balb akr non nod lp
 142 155 130 167 142 138 142 167 138 142 138 167

MTH227 D8Mit148

Thu 4 Nov 1993

MAP POSITION: 64.9 cM from top of Chr8 (Framework)

PRIMERS: CATCTCCCCCTCACCTCATA

TGTTAGCAATATGGATTTTAGAGCC

PRODUCT: 238 bp

GENOTYPE: HBABBAHHAHBAHHHHHHBAAAB BAHABBABBBHHHAHBHHHHHH (rank 1)

SEQUENCE:	10	20	30	40	50	60
0	ACACCCACAT	CTCCCCCTCA	CCTCATATCT	ACACCTACAC	ACACACACAC	ACACACCTGC
60	ACATGTACAC	ACACACACCT	GCACATGTAC	ACACACACAC	ACACACACCT	GCACATGTAC
120	ACACACACAC	ACACACACAC	ACACCTGTAC	ATGTACACAC	ATAGACACAC	ACACACACAC
180	ACACACACAC	ACACACACAC	TTNGTAGATC	TTGAGTATGA	GGCTCTAAAA	TCCATATTGC
240	TAACAGGAGA	TAGCTCTG				

ALLELES: ob cast spr a b6 c3h dba balb akr non nod lp
 239 227 369 239 239 243 239 239 243 239 239 227

M77 D8Mit13

Tue 27 Apr 1993

MAP POSITION: 64.9 cM from top of Chr8 (Framework)

PRIMERS: CCTCTCCAGCCCTGTAAG

AACGTTTGTGCTAAGTGGCC

PRODUCT: 98 bp

GENOTYPE: HBABBAHHAHBAHHHHHHBAAB BAHABBABBBHHHAHBHHHHHH (rank 1)

SEQUENCE:	10	20	30	40	50	60
0	TTNAAGCNG	GTCCCGGGGA	TCTCATTATG	GATGGNTGTG	AGCCACCATG	TGTAGGCTGG
60	GAATTGAACT	CAGGACCTTT	GGAAGAGCAG	TCAGTGCTCT	TAACCACTGA	GCCTCTCTCC
120	AGCCCTGTAA	GAATATTTCT	TACACACACA	CACACACACA	CACACACACA	CACACGGAGG
180	GGGAGAAATG	GCCACTTAGC	ACAAACGTTG	ATAAAGCAGA	GGCTACACTA	CGTACTACAC
240	GGGCATAACC	ACGGCACCCA	CAGGATGGAC	GGTGATTATC	ACAAAGCTCT	TACACTTCAC
300	AGGAAGTGTG	GGCTGTGATG	ATGCAGAGAC	CTATGCTAGC	TTCCCAGAGC	CACCAACAAA
360	TGATGGACCT	CACAAGACGT	TATTATAGAG	AAGCGCCAAA	AGGNACCACC	CAGAT

ALLELES: ob cast spr a b6 c3h dba balb akr non nod lp
 98 114 114 98 98 91 98 105 94 98 98 108

^L

MAP BASE QUERY REPORT

129 is like this

Bill Dietrich
67) 252 1915

Thurs. June 2nd

Genotyping 2/1 v. 2/2

Kim has ~~been~~ started for any markers in Aqt gene & none found. Squire & Maelene tested quantitative Southern for genotyping 2/1 v. 2/2. Very unsatisfactory. Better logic is that in F2 the heterozygotes are always B6 and 129-derived, the homozygotes are B6 (normal) or 129 (doubly-mutant). Any marker closely linked to Aqt can be used for genotyping. First tests with D8 Mit 13 and Mit 14 done by Kim look promising (at 64.9 cM). Aqt is at ~70 cM. D8 Mit 156 is at 70.5 cM & so c'd be better in theory. Test D8 Mit 156 and 56, using primers to left. Note lp is fairly close to 129 in Fetch's genealogy, & 13 & 14 give predicted (lp) sizes.


Mon 5 Oct 1992

B740 D8Mit56
MAP POSITION: 70.5 cM from top of Chr8 (Placement)
PRIMERS: ACACCTCAGAGACCATGAGTACACC GAGTTCACCTACCCACAAGTCTCC
PRODUCT: 162 bp
GENOTYPE: HBAHHAAHBAHHHHAHBAAH- BABABBABBBHHHAABHHHHHH (rank 1)
SEQUENCE: 10 20 30 40 50 60
0 CCGCTCCTGA GACTACTTCA GACTCTCAGA GACCATGAGT ACACCACTAA GAGGAGCAAG
60 TAAATGGTGG AGAAACAGAA GAAAAAGAGA AACAGGAGGA TATGGGCGCG CACAAACACA
120 CACACACACA CACACACACA CCGCATAACT GGAGACTTGT GGAGACTTGT GGGTAGTGAA
180 CTCAATTAAT TAGGCTTAAT TTGAGTAGTT
ALLELES: ob cast spr a b6 c3h dba balb akr non nod lp
160 184 152 164 160 164 182 164 164 160 180 182
MAP BASE QUERY REPORT

Thu Jun 2 13:50:25 EDT

Mon 15 Nov 1993

MT1329 D8Mit156
MAP POSITION: 70.5 cM from top of Chr8 (Placement)
PRIMERS: ATTCTGCAATGCAAATACATACA AACCAATGGCCTCCAGAAG
PRODUCT: 139 bp
GENOTYPE: HBAHHAAHBAHH-HAHBAHBAH BABABBABBBHHHAABHHHHHH (rank 1)
SEQUENCE: 10 20 30 40 50 60
0 ATTCTGCAAT GCAAATACAT ACACACACAC ACATACACAC ACACACACAC ACAGAGAGAG
60 AGAGAGAGAG AGAGAGAGAG AGAGAGAAT TGTGGTTTCA CTGACCTTTC AAGTGGTGGC
120 CTCTGGAGG CCATTGGTTC CAGTCACAAA TATAGATACT ATTGTGATAA GCAAGTTAAA
180 TACAAATGAG TACCCCAAGA GCCTAAGAAG TTATCTTGTT AGAACCCTCA GGNCTCCTTT
240 TATTTATGGC ATCTAGTATA GGAAAGNCTC TCAACTAACC TGAGAAGTAG GTCTCAAGAA
300 ACATGCTGGT ATCAAGGAAA TGAGTGCAGA TGTATGAGTA AT
ALLELES: ob cast spr a b6 c3h dba balb akr non nod lp
140 158 150 154 140 154 154 154 154 140 154 152



The chart shows a current genetic map of the mouse that provides a marker every 0.6 centimorgans (cM) on average, which is roughly equivalent to every 1.1 megabases. The genetic markers are of two types: 1518 simple sequence length polymorphisms (SSLPs), shown on the left, and 1098 gene-based loci, shown on the right.


SSLP Map. The SSLP map consists of loci defined by polymerase chain reaction (PCR) assays, each involving a specific pair of PCR primers flanking the site of a short repeat sequence of variable length — in most cases, (CA)_n. These markers tend to be highly polymorphic among inbred laboratory strains, making them especially useful for genotyping intraspecies crosses. Most were based on anonymous (random) sequences, although about one-tenth were taken from repeats found in known genes. The SSLP markers shown were all genotyped in a single interspecies F₂ cross [W. Dietrich *et al.*, *Genetics*, 131,423 (1992)]. The data were subjected to a mathematical error-checking procedure [(S.E. Lincoln and E.S. Lander, *Genomics* 14, 604 (1992))], with the result that the relative positions of the loci were supported by a likelihood ratio of 1000:1, except for underlined loci, for which the ratio was >10:1. For nearly 98% of the markers, which were developed at the Whitehead Institute/MIT Center for Genome Research (WI-CGR), Cambridge, MA, formal locus names have been abbreviated. For example, D17Mit3 is shown as 3 on chromosome 17. The remaining SSLPs were developed elsewhere but were also genotyped in the same cross. For these loci, the symbol of the originating laboratory was retained (For example, D4Nds1 is abbreviated Nds1).

Gene-Based Map. The gene-based map is primarily defined by cloned gene probes that detect restriction fragment length polymorphisms (RFLPs). This map is especially valuable for comparative mapping between the mouse and human genomes. The framework for the map is the Frederick interspecies backcross map [N.G. Copeland and N.A. Jenkins, *Trends Genet.* 7,113 (1991)], shown in *black type*. Because these loci were all mapped in a single backcross and subjected to mathematical error-checking, the orders of loci were supported by a likelihood ratio of 1000:1, except for underlined loci for which the ratio was >10:1. Loci shown in *red* (■) consist of additional genes and anonymous DNA segments mapped in human and mouse that were not mapped in Frederick, but were reported in the 1993 Mouse Chromosome Committee Reports (*Mammalian Genome*, in press); the positions of these genes were inferred on the basis of mapping information in these committee reports. Loci in brackets <> were used to align the maps with respect to the centromere. The position of the centromere is indicated by a circle at the top of each chromosome. Chromosome lengths were drawn to scale on the basis of a 1600-cM genetic length. A break in the distal region of chromosome 11 indicates that this chromosome is slightly longer than predicted by a 1600-cM map.

Integration of Maps. The two genetic maps were integrated by mapping approximately 250 of the SSLPs from the WI-CGR in the Frederick interspecies backcross. These SSLPs are indicated in *green* (■), with lines connecting their locations in the two crosses.

Mouse-Human Homology. The colored segments within each chromosome map indicate known regions of syntenic or linkage conservation between the mouse and human genomes. The correspondence is based on those mouse genes whose human homologs have been mapped in the human genome; these loci are indicated in *bold*. For loci mapping near the boundary of two human homology segments, those loci mapping to the proximal segment are marked with an asterisk.


Table. The table included in the chart contains a representative list of



Fre. Sun

MIT 13/14

Kim tested 13 and 14 -
Recombinants will arise, do
from Agt. Recombinant do

→ 129 
These have no effect on type

129 


↑
This will be a 1 chromosome
but will M13 type as 129


+ 

This animal is 1/1
but will M13 type
as 2/1. No problem
because it will
be Neo⁻
(will measure map
distance)

+ 

This animal is 2/1
but will M13 type
as 2/2. It
will ∴ score
incorrectly.



+ 


This animal is 2/1
but will M13 type
as 1/1. No problem
because it will
be Neo⁺
(will measure map
distance)


9181
maybe
such

+ 

This animal is 2/2
but will M13 type
as 2/1. It
will ∴ score
incorrectly

So 50% of recombinants will be recognized & cause no error
Other 50% will not be recognized & will blur the edges of 2/1 & 2/2





MIT 56 looks good, also
but 56 is not good
(c.f. mp. of primer)

Back - CR P2 F2 F1 129 F2 129 F2 F1 86 C3H
SS99 SS99 8762 EL 9182 mix 9181 9468 9310 MIT 56





Fig. 1. (A) Genealogy of 24 inbred strains of mice (10-12). Black dots refer to crosses made in production of a particular strain. (B) Parsimony analysis of 144 single locus genotypes (rather than a genetic distance matrix) for 24 inbred strains of mice. Phylogenetic trees representing estimated relationships among strains were produced by using the parsimony algorithm (15). The length of the tree (L) reflects a total of 565 allele fixations. The branch lengths are in allele fixations (arabic numbers). The numbers in italics reflect the number (out of 100 resamplings of the data) that give the same clade (16). Nonintegral numbers of substitutions (allele fixations) arise from averaging over all possible parsimonious solutions for the tree. Dotted lines represent alternative but equally parsimonious joinings of taxa.



Resting / Conscious

Data on crossing over (absence)

2/1 2/2 total

37-120	2	3	5
37-112	6	5	11
37-94	7	3	10
37-76	15	6	21
37-66	22	6	28
37-63	29	10	39
37-58	10	13	23

91 46 137

B6/129 129/129

Mit 56.

Absence of any Neo⁺ B6⁺
129 shows no X.O. in
v91 meioses.

11/25/94

AGT Neo AGT 56 Long (129)
AGT 56 Short (B6)

(2)
(1)

Recombinant Platelet-derived Growth Factor B Gene Expression in Porcine Arteries Induces Intimal Hyperplasia In Vivo

Elizabeth G. Nabel,* Zhiyong Yang,* Susanne Liptay,* Hong San,* David Gordon,* Christian C. Haudenschield,* and Gary J. Nabel*

*Departments of Internal Medicine, Biological Chemistry, and Pathology, Howard Hughes Medical Institute, University of Michigan, Ann Arbor, Michigan 48109; and †Jerome H. Holland Laboratory, American Red Cross, Rockville, Maryland 20855

Abstract

Platelet-derived growth factor (PDGF) B chain induces cell proliferation in vitro and is associated with arterial lesions that cause cardiovascular disease. However, it has been difficult to document the biological response to PDGF B gene expression in arteries in vivo. To determine the biologic effects of this growth factor in vivo, we have introduced an eukaryotic expression vector plasmid encoding recombinant PDGF B by direct gene transfer into porcine iliofemoral arteries using DNA liposome complexes. The presence of PDGF B plasmid DNA and expression of recombinant mRNA were confirmed by polymerase chain reaction analysis, and recombinant PDGF protein was demonstrated by immunohistochemistry. Intimal thickening was observed in porcine arteries 21 days following transfection with the recombinant PDGF B gene compared with arteries transduced with a control gene, *E. coli* β -galactosidase. An eightfold increase in intimal to medial ratio was present in PDGF B gene transfected arteries compared with control transfected arteries ($P = 0.001$). This study suggests that expression of a recombinant PDGF B gene in vivo can play a role in the induction of intimal hyperplasia, which can lead to cardiovascular diseases. (*J. Clin. Invest.* 1993. 91:1822-1829.) Key words: gene transfer • gene expression • platelet-derived growth factor • cellular proliferation • liposomes

Introduction

Multiple growth factors stimulate vascular smooth muscle cell proliferation in vitro, including the dimeric forms of PDGF. PDGF BB is a potent smooth muscle cell mitogen (1, 2) that may also act as a smooth muscle chemoattractant (3-5). In vivo, the role of the PDGF B gene in tissue injury and atherosclerosis has been difficult to address. This difficulty arises from the complexity of cellular and protein interactions in

vivo, their paracrine and autocrine nature, and the cascade of events that can be induced by experimental manipulations. It is often difficult to determine whether a specific gene product induces intimal hyperplasia directly or indirectly through other gene products synthesized in response to injury. Recently, it has become possible to express recombinant genes in vivo by direct gene transfer utilizing retroviral vectors or DNA liposome complexes (6-9). Direct gene transfer permits the delivery of a specific recombinant gene into vascular cells at specific sites in vivo and affords the opportunity to determine the effects of a specific gene product in the arterial wall (10). To define the function of PDGF B within arteries, we have directly introduced the recombinant gene in porcine arteries in vivo and have shown that expression of recombinant PDGF B can cause intimal hyperplasia. These data suggest that expression of PDGF B can induce vascular lesions relevant to the pathogenesis of atherosclerosis and restenosis.

Methods

Plasmids and cell transfection. The PDGF B expression vector used for in vitro and in vivo experiments was prepared by ligating the SalI-XbaI fragment of the v-sis gene into the XhoI-BamHI cloning site of the pSVL vector (11) (Pharmacia, LKB Biotechnology, Piscataway, NJ). This vector utilizes the SV40 late promoter to regulate expression of the PDGF B gene. Primary porcine endothelial cell cultures were established as previously described (12) to test expression of the PDGF B vector. The pSVL/PDGF B plasmid expression vector was transfected into porcine endothelial cells using 12 μ g of Lipofectin® (Bethesda Research Laboratories, Gaithersburg, MD) and 5 μ g of DNA (stock concentration > 1 mg/ml) (13). Culture transduced endothelial cells were assayed for secretion of recombinant PDGF BB supernatants (10%) with a colorimetric proliferation assay (14) using NIH 3T3 cells, and a fourfold or greater increase in proliferative activity was observed. No evidence of cell transformation was observed in endothelial or vascular smooth muscle cells stably transduced with this vector. The *E. coli* β -galactosidase expression vector was prepared by cloning the β -galactosidase gene into the pDOL vector, which was derived from the Moloney murine leukemia virus (Mo-MuLV) (15). The wild type Mo-MuLV LTR provided the promoter for the β -galactosidase gene.

Gene transfer in vivo. In vivo arterial gene transfer was performed in 15 pigs, 9 with the recombinant PDGF B gene and 6 with a control reporter gene, *E. coli* β -galactosidase. Double balloon catheters were positioned in the iliofemoral arteries as previously described (6). The arterial segment was rinsed with 5 ml saline and 5 ml Opti-MEM to clear the vessel of blood. Approximately 10 min before the insertion of the catheter, the DNA liposome conjugates were prepared. 5 μ l lipofectin was diluted into 0.2 ml Opti-MEM at room temperature, and 2-5 μ g plasmid DNA (stock concentration > 1 mg/ml) was added and mixed by gentle tapping. The solution remained at room temperature for 5-10 min, and 0.5 ml Opti-MEM was added to the DNA liposome

Send correspondence and reprint requests to Elizabeth G. Nabel, M.D., University of Michigan, MSRB II, Room 3560, 1150 W. Medical Center Drive, Ann Arbor, MI 48109-0688.

Received for publication 17 July 1992 and in revised form 23 December 1992.

J. Clin. Invest.

© The American Society for Clinical Investigation, Inc.

0021-9738/93/04/1822/08 \$2.00

Volume 91, April 1993, 1822-1829

Mon. June 20th

On atherosclerotic plaques

N.M.'s data on log distribution of plaque sizes, combined with published work on clonal nature of atherosclerotic plaques, suggests, as published, that they are partially transformed (c.f. anaplastic lesions, pre-cancer). If so they should grow better in culture.
 • Next with Ed Sheehy, the growth of $F2$ E^+/E^+ versus E^-/E^- vascular endothelial cell cultures.

Wed. June 22nd

Nobuyo comments that the (transformed) cells are first macrophages, then altered muscle cells, so we may have to look for strange outgrowths -

O.S. - anyway transplantability is what I'm looking for at first.

Thurs. June 23rd (O. & R.'s birthday 69!)

~~BGF~~ B c'd be a candidate (see Nobel et al. J.C.I. 91 1822 (1993) - brought by Bob Redfiek)

Fri. June 24th

- on its receptor!

Mon. June 24th

With Nobunjo

$$V = \frac{V_{\max} [S]}{K_m + [S]}$$

$$\therefore \frac{V}{V_{\max}} = \frac{[S]/S_{0.5}}{1 + [S]/S_{0.5}} \quad (S_{0.5} = K_m)$$

$$\text{or } y = \frac{x}{1+x}$$

$$\therefore \frac{dy}{dx} = \frac{1}{1+x} - \frac{x}{(1+x)^2}$$

$$\text{As } x \rightarrow 0 \quad \frac{dy}{dx} \rightarrow 1$$

$$\text{as } x \rightarrow \infty \quad \frac{dy}{dx} \rightarrow 0$$

$$\text{At } x=1 \quad \frac{dy}{dx} = \frac{1}{2} - \frac{1}{4} = \frac{1}{4}$$

$\therefore$ slope at $[S] = K_m$ is $4 \times$ slope
at $[S] = K_m/100$

Human

Mouse

Hence mo. is $\sim 4 \times$ more sensitive
to $[AGT]$ than human

Shingles!!



Let's see what happens when the clearance varies
(receptor)

$$2p = \text{Utilization} + \text{Clearance} = \text{Constant}$$

clearance capability

$\therefore$ if ~~clearance~~ decreases the concentration of AGT will $\uparrow$ until new utilization & new clearance are at equilibrium.

Consider a heterozygote for disruption of clearance gene. Let new level of AGT be $r \times$ old.

$\leftarrow$ Assume util. is $\propto [AGT]$

$$\text{Then } U_{100} + C_{100} = r \cdot U_{100} + r \cdot \frac{C_{100}}{2}$$

$$\text{But } U_{100} = 0.3 \times C_{100}$$

$$\therefore 1.3C_{100} = r(0.3 + 0.5)C_{100}$$

$$\text{And } r = \frac{1.3}{0.8} = 1.63$$

Thus we can predict that $[AGT]$ will increase 1.63 fold if clearance of AGT decreases to half.

This type of reasoning will apply to the ANP clearance receptor. It may apply to the RIA and RIB experiments, since the clearance of ATII will be changed, but less likely; more likely r_{in} will $\uparrow$ & $[AGT]$ will go down and ATII will $\uparrow$.

Sunday July 31

Same more equations (see B'78)

let's see what solutions follow if utilⁿ of AGT $\rightarrow$ ATT
& thence bp. $\uparrow$ with $\uparrow$ in gene copy.

$\therefore$ let U (utilization) be proportional to steady state
bp, with bp = $\frac{\text{sum}}{\text{num}}$ in the 1/1 animal and with
sum Hg change / copy number

Then

$$1/1 \quad 2p = U_{100} + C_{100} \quad (1)$$

$$1/0 \quad p = 0.92 U_{100} + 0.35 C_{100} \quad (2)$$

$$2/1 \quad 2px + p = 1.08 U_{100} + 1.24 C_{100} \quad (3)$$

substituting (2) into (1) gives

$$1.84 U_{100} + 0.7 C_{100} = U_{100} + C_{100}$$

$$\text{or } 0.84 U_{100} = 0.3 C_{100}$$

$$\text{and } U_{100} = 0.357 C_{100} \quad \text{Was } 0.3$$

$$\text{And } p = \frac{1.357 C_{100}}{2} = 0.679 C_{100} \quad \text{Was } 0.65$$

$$\begin{aligned} \text{And } 2px &= 1.08 \times 0.357 C_{100} + 1.24 C_{100} - 0.679 C_{100} \\ &\quad (0.386) \\ &= 0.947 C_{100} \end{aligned}$$

$$x = \frac{0.947 C_{100}}{2 \times 0.679 C_{100}} = 0.697 \quad \text{Was } 0.685$$

This variable utilization does not
render the system invalid.

(Cont.)

if not, will it be purified later by ...

5' 08277 note: CR 5-13 1LEFT

25
52%

G	C	T	C	A	A	A	A	G	A	T	C	G	A	G
G	T	G	C	T	T	G	T	G	C					

5' 08278 note: CR 140-147 1MIDLEFT

23
52%

G	A	G	A	T	G	A	T	G	C	T	C	G	C	T
C	T	G	T	T	T	C	G							

5' 08279 note: CR 148-141 1MIDRIGHT

23
52%

T	G	G	C	G	A	A	A	C	A	G	A	G	C	G
A	G	C	A	T	C	A	T							

08280 5' 1Right note: CR 209-140 ← mislabelled
sh'd be 202

23
52%

C	T	G	C	C	T	T	G	G	A	T	G	T	A	G
C	G	C	A	C	T	A	T							

Overall will be 613 bp 5' portion will be 427 bp 3' will be 200 bp

A different "middle" could be in the rat territory

(+125)

E Y S H L T R V A P

or better A(109)

M L S A G A L A A G F

— but try no. sequences first.

App of work


Tues. Aug. 2nd

ANP Clearance Receptor (CR)



In view of importance of clearance in modulating concentrations, I will try to alter $\uparrow$ & $\downarrow$, the ANP Clearance Receptor.

Sequence data on Human ANP Receptor (Type 1)
from Sahygi et al. J. BC. 266 1122 (1991) &
from Fuller et al. J. B. C. 263 9395 (1988) &
on rat & mouse (cDNA) from David Lowe (unpublished)

Structure: —



Since the 3' end is of indeterminate length, it is probably best to keep 5' short & 3' long. The 5' clone is easier for a beginning.



6 clones 211 w'd be > 600bp or 2 x ~ 300 for 2 probes for 1 probe

All these worked
S1/III
w/16

5' ACC ^{promoter} probe was ~1 kb
hatched on 3' ACC was ~480 bp

w/1 5' ACC ^{intron 2} probe was ~1 kb
488 3' UT 3' ACC was ~270 bp (cut off)

labelled B'108 & etc. Taken up in 500 μ l TE 1X CIAA

Summary of these first primer pairs

Full name	Trivial Name	Expected length	Comments
CR 5-13	<u>1 Left</u> (exon 1 5')	615 bp	All coding sequence of mature protein (whole protein) exon 1
CR 209-140	<u>1 Right</u> (exon 1 3')		
CR 5-13	<u>1 Left</u>	434 bp	Uses the first available mouse data it is first 2/3 of exon 1
CR 148-141	<u>1 Mid right</u> (mid exon 1 mouse data)		
CR 140-147	<u>1 Mid left</u> (mid exon 1 mouse data)	210 bp	mouse data only it is the second 1/3 of exon 1
CR 209-140 ↑ slide for 202	<u>1 Right</u> (exon 1 3')		

Test with all three. Hope to use the 2/3 & 1/3 for double lifts

CR 433-440	<u>7 Left</u> (exon 7 5' end)	Uncertain - prob. $\sim$ 1 kb	Exon 7 Intron 8 Exon 8
CR 488-481	<u>8 Right</u> (exon 8 3' end)		

Wed. Aug 3rd

3' end of ANP Clearance Receptor (CR)

The 3' end is complicated by the presence of an 8
3' untranslated region of unknown length, but

08286	5'	note: CR488-481										8 Right	3'			
le: -		T	C	T	T	C	T	C	G	C	A	G	C	T	C	T
l: 2.3		C	G	A	T	G	C	T	T							
52%																

08287	5'	note: CR433-440										7 left	3'			
le: -		G	C	C	T	A	G	A	A	G	A	A	T	C	T	G
l: 2.4		C	A	G	T	G	A	C	A	G						
50%																

length 25 GC 44% ← Not v. good

or better

G	K	H	R	E	L	R	E	D	S
GGG	AAG	CAT	CGA	GAG	CTG	CGA	GAA	GAT	TCC
110	910	921	910	940	901	911	911	911	911

length 23 GC 52% CR 488-481

433 440

G	L	E	E	S	A	V	T	G	I	V
GGC	CTA	GAA	GAA	TCT	GCA	GTG	ACA	GGA	ATC	GTT

length 24 GC 50% CR 433-440

header

Irind Rpd
Primer
Irind best

1
Power
1 Right

2017

8 Right

From David
Howe


-40	rNPRC	M	R	S	L	L	F	T	F	S	A	C	V	L	A	R	A	L	L	A	G	G	A	-	-	-	-	S	S	G	G	D	I	G	P	G	N	R	R	E	A	L	A								
-45	hNPRC	H	P	S	L	V	L	T	F	S	P	C	V	L	G	W	A	L	L	A	G	G	T	G	G	G	G	G	G	G	G	G	G	G	G	G	G	G	R	Q	E	R	E	A	L	P					
-41	bNPRC	M	P	S	L	V	L	T	F	S	A	C	V	L	L	G	W	A	L	L	A	D	C	T	G	-	-	-	G	G	G	S	G	G	A	G	P	G	R	G	R	R	E	A	L	P	P				
6	rNPRC	Q	K	I	E	V	L	V	L	P	R	D	S	Y	L	F	S	L	A	R	V	R	P	A	I	E	Y	A	L	R	S	V	E	G	N	G	T	G	R	K	L	L	P	P	G	T	R	F	O		
6	hNPRC	Q	K	I	E	V	L	V	L	P	O	D	S	Y	L	F	S	L	T	R	V	R	P	A	I	E	Y	A	L	R	S	V	E	G	N	G	T	G	R	L	L	P	P	G	T	R	F	O			
6	bNPRC	Q	K	I	E	V	L	V	L	P	O	D	S	Y	L	F	S	L	A	R	V	R	P	A	I	E	Y	A	L	R	T	V	E	G	N	A	T	G	R	R	L	L	P	A	G	T	R	F	O		
56	rNPRC	V	A	Y	E	D	S	D	C	G	N	R	A	L	F	S	L	V	D	R	V	A	A	A	G	A	K	P	D	L	I	L	G	P	V	C	E	Y	A	A	A	P	V	A	R	L	A	S	H	W	
56	hNPRC	V	A	Y	E	D	S	D	C	G	N	R	A	L	F	S	L	V	D	R	V	A	A	A	G	A	K	P	D	L	I	L	G	P	V	C	E	Y	A	A	A	P	V	A	R	L	A	S	H	W	
56	bNPRC	V	A	Y	E	D	S	D	C	G	N	R	A	L	F	S	L	V	D	R	V	A	A	A	G	A	K	P	D	L	I	L	G	P	V	C	E	Y	A	A	A	P	V	A	R	L	A	S	H	W	
106	rNPRC	D	L	P	M	L	S	A	G	A	L	A	A	G	F	O	H	K	D	T	E	Y	S	H	L	T	R	V	A	P	A	Y	A	K	M	G	E	M	L	A	L	F	R	H	H	W	S	R	A		
106	hNPRC	D	L	P	M	L	S	A	G	A	L	A	A	G	F	O	H	K	D	S	J	E	Y	S	H	L	T	R	V	A	P	A	Y	A	K	M	G	E	M	L	A	L	F	R	H	H	W	S	R	A	
106	bNPRC	D	L	P	M	L	S	A	G	A	L	A	A	G	F	O	H	K	D	T	E	Y	S	H	L	T	R	V	A	P	S	Y	A	K	M	G	E	M	L	A	L	F	R	H	H	W	S	R	A		
156	rNPRC	A	L	L	Y	S	D	D	K	L	E	R	N	C	Y	F	T	L	E	G	V	H	E	V	F	O	E	E	G	L	H	T	S	A	Y	N	F	D	E	T	K	D	L	D	I	D	I	V	R	Y	
156	hNPRC	A	L	V	Y	S	D	D	K	L	E	R	N	C	Y	F	T	L	E	G	V	H	E	V	F	O	E	E	G	L	H	T	S	I	Y	S	I	F	D	E	T	K	D	L	D	E	I	V	R	N	
156	bNPRC	V	L	V	Y	S	D	D	K	L	E	R	N	C	F	F	T	L	E	G	V	H	E	V	F	O	E	E	G	L	H	T	S	A	Y	N	F	D	E	T	K	D	L	D	E	I	V	R	H		
206	rNPRC	I	Q	G	S	E	R	V	I	M	C	A	S	D	I	R	I	R	I	M	L	A	V	H	R	H	G	M	T	S	G	D	Y	A	F	F	N	I	E	L	F	N	S	S	Y	G	D	G			
206	hNPRC	I	Q	A	S	E	R	V	I	M	C	A	S	D	I	R	S	I	M	L	V	A	H	R	H	G	M	T	S	G	D	Y	A	F	F	N	I	E	L	F	N	S	S	Y	G	D	G				
206	bNPRC	I	Q	A	S	E	R	V	I	M	C	A	S	D	I	R	G	I	M	L	A	A	H	R	H	G	M	T	S	G	D	Y	A	F	F	N	I	E	L	F	N	S	S	Y	F	Y	G	D	G		
256	rNPRC	S	W	K	R	G	D	K	H	D	F	E	A	K	Q	A	Y	S	S	L	O	T	V	T	L	L	R	T	A	K	P	E	F	E	K	F	S	M	E	V	K	S	S	V	E	K	O	G	L	N	
256	hNPRC	S	W	K	R	G	D	K	H	D	F	E	A	K	Q	A	Y	S	S	L	O	T	V	T	L	L	R	T	Y	K	P	E	F	E	K	F	S	M	E	V	K	S	S	V	E	K	O	G	L	N	
256	bNPRC	S	W	K	R	G	D	K	H	D	F	E	A	K	Q	A	Y	S	S	L	O	T	I	L	L	R	T	A	K	P	E	F	E	K	F	S	M	E	V	K	S	S	V	E	K	O	G	L	S		
306	rNPRC	E	D	Y	V	N	M	F	V	E	G	F	H	D	A	I	L	L	Y	V	L	A	H	E	V	L	R	A	G	Y	S	K	K	D	G	G	K	I	I	Q	O	T	W	N	R	T	F	E	I		
306	hNPRC	E	D	Y	V	N	M	F	V	E	G	F	H	D	A	I	L	L	Y	V	L	A	H	E	V	L	R	A	G	Y	S	K	K	D	G	G	K	I	I	Q	O	T	W	N	R	T	F	E	I		
306	bNPRC	E	D	Y	V	N	M	F	V	E	G	F	H	D	A	I	L	L	Y	V	L	A	R	E	V	L	R	A	G	Y	S	K	K	D	G	G	K	I	I	Q	O	T	W	N	R	T	F	E	I		
356	rNPRC	A	G	O	V	S	I	D	A	N	G	D	R	Y	G	D	F	S	V	I	A	M	T	D	E	A	G	T	O	E	V	I	G	D	Y	F	G	K	E	G	R	F	K	M	R	S	N	V	K	Y	
356	hNPRC	A	G	O	V	S	I	D	A	N	G	D	R	Y	G	D	F	S	V	I	A	M	T	D	V	E	A	G	T	O	E	V	I	G	D	Y	F	G	K	E	G	R	F	E	M	R	P	N	V	K	Y
356	bNPRC	A	G	O	V	S	I	D	A	N	G	D	R	Y	G	D	F	S	V	I	A	M	T	D	E	A	G	T	O	E	V	I	G	D	Y	F	G	K	E	G	R	F	E	M	R	P	N	V	K	Y	
406	rNPRC	P	W	G	S	L	K	L	R	I	D	E	T	R	I	V	E	H	T	N	S	P	C	K	S	C	-	G	L	E	E	S	A	V	T	G	I	V	V	G	A	L	L	G	A	G	L	L	M	A	
406	hNPRC	P	W	G	P	L	K	L	R	I	D	E	N	R	I	V	E	H	T	N	S	P	C	K	S	C	-	G	L	E	E	S	A	V	T	G	I	V	V	G	A	L	L	G	A	G	L	L	M	A	
406	bNPRC	P	W	G	P	L	K	L	R	I	D	E	T	R	M	V	E	H	T	N	S	P	C	K	A	S	G	L	E	E	S	A	V	T	G	I	V	V	G	A	L	L	G	A	G	L	L	M	A		
455	rNPRC	F	Y	F	F	R	K	K	Y	R	I	T	I	E	R	R	N	H	Q	E	E	S	N	I	G	K	H	R	E	L	R	E	D	S	I	R	S	H	F	S	V	A									
455	hNPRC	F	Y	F	F	R	K	K	Y	R	I	T	I	E	R	R	T	Q	Q	E	E	S	N	L	G	K	H	R	E	L	R	E	D	S	I	R	S	H	F	S	V	A									
456	bNPRC	F	Y	F	F	R	K	K	Y	R	I	T	I	E	R	R	N	O	Q	E	E	S	N	V	G	K	H	R	E	L	R	E	D	S	I	R	S	H	F	S	V	A									

ratNPRC 1 GAATTCGGCTTGCTGGCAACAGGATGCGGTCCCTGCTGCTGTTCACTTTC
ratNPRC 51 TCGGCGTGCGTGCTGCTGGCCCGGGCGCTGCTGGCTGGCGGTGCGAGTAG
ratNPRC 101 TGGCGGTGGGGACACCGGGCCAGGCAACAGGCGCGGGAGAGAGAGGCGC
ratNPRC 151 TGGCGGCTCAAAAGATCGAGGTGCTTGTGCTATTGCCCGGGACGATTTCG
ratNPRC 201 TACTTGTTCTCGCTGGCCCGGGTGAGGCCGGCCATCGAGTACGCGCTGCG
ratNPRC 251 TAGCGTGGAGGGCAATGGCACCGGCAGGAAGCTGCTGCCCGCCGGCACTC
ratNPRC 301 GCTTCCAGGTGGCCTATGAAGACTCTGACTGTGGCAACCGTGCGCTCTTC
ratNPRC 351 AGTTTGGTGGACCGCGTGGCGGGCGCGCGCGGCCAAGCCGGACCTCAT
ratNPRC 401 CCTGGGTCCGGTGTGCGAGTACGCTGCGGGCGCCGGTGGCCCGGCTGGCGT
ratNPRC 451 CTCAGTGGGACCTGCCGATGCTGTCCGCAGGAGCGCTGGCCCGGGGTTTC
ratNPRC 501 CAGCACAAGGACACGGAGTACTCGCACCTCACGGCGTGGCGCTGCCTA
ratNPRC 551 CGCCAAGATGGGAGAGATGATGCTCGCTCTGTTCCGCCACCACTGGA
mNPRC 1 A K M G GAGATGATGCTCGCTCTGTTCCGCCACCACTGGA
ratNPRC 601 GCCGCGCTGCCGTGCTCTACAGCGACGACAAACTCGAGAGAAACTGCTAC
mNPRC 38 GCCGTGCA GCCCTGTCTCTACAGCGACGACAAACTCGAGAGGAACTGTTAT
ratNPRC 651 TTCACCTCTCGAGGGGGTCCACGAGGTTTTCCAGGAGGAGGGCTTGACAC
mNPRC 88 TTCACCTCTCGAGGGGGTCCACGAGGTTTTCCAGGAGGAGGGCTTGACAC
ratNPRC 701 GTCTGCCTACAATTTTCGACGAGACCAAAGACTTGGACCTGGACGACATAG
mNPRC 138 GTCTGCCTACAATTTTCGACGAGACCAAAGACTTGGACCTGGACGACATAG
ratNPRC 751 TGGCGCTACATCCAGGGCAGTGAGCGAGTGGTGATCATGTGTGCCAGTGGT
mNPRC 188 TGGCGCTACATCCAGGGCAGTGAGCGAGTGGTGATCATGTGTGCCAGTGGT
ratNPRC 801 GACACCATTCCGAGAATCATGTGGCGGTGCACAGACACGGCATGACCAG
mNPRC 238 GACACCATTCCGAGAATCATGTGGCGGTGCACAGACACGGCATGACCAG
ratNPRC 851 TGGAGACTATGCTTTCTTCAACATTGAACCTCTTCAACAGTTCTCTTACG
mNPRC 288 TGGAGACTAGCTTTCTTCAACATTGAACCTCTTCAACAGTTCTCTTACG
ratNPRC 901 GAGATGGCTCGTGGAGAGAGGGGACAAACACGACTTGAAGCTAAGCAA
mNPRC 338 GAGATGGCTCGTGGAGAGAGGGGACAAACACGACTTGAAGCTAAGCAA
ratNPRC 951 GCTTACTCATCCCTCCAAACAGTCACCCTGCTGAGGACCGGGAAGCCTGA
mNPRC 388 GCTTACTCATCCCTCCAAACAGTCACCCTGCTGAGGACCGGGAAGCCTGA
ratNPRC 1001 GTTTGAGAAAGTTTTCCATGGAGGTGAAAAGTTCTGTTGAGAAACAAGGGC
mNPRC 438 GTTTGAGAAAGTTTTCCATGGAGGTGAAAAGTTCTGTTGAGAAACAAGGGC
ratNPRC 1051 TCAATGAGGAGGATTACGTGAACATGTTGTTGAAGGCTTCCATGATGCC
mNPRC 488 TCAATGAGGAGGATTACGTGAACATGTTGTTGAAGGCTTCCATGATGCC
ratNPRC 1101 ATCCTCCTCTACGTTCTGGCTTTCATGAAGTCTCAGAGCTGGCTACAG
mNPRC 538 ATCCTCCTCTACGTTCTGGCTTTCATGAAGTCTCAGAGCTGGCTACAG

Mature Protein
1L2

IMR2

300



ON 7.15 pm

OFF 7.49 pm

10V/cm

Bye 1.5% Agarose
W Hx1

A (undiluted) 1L 1R

B " 1L 1MR

C " 1ML 1R

W

Acryl. marker

W

Bye

Conclusion: The mouse pair, C, worked.

The rat sequence L1, is no good
Purify the 210bp, 3' end of exon 1, probe
- & try Southern etc.

Thurs. Aug 8th

ANP - CR probes

PCR formulation from A'121 for a ~~25~~ ²⁵ ~~pl~~ RX

X 226 μ l
✓ 17.5 μ l
✓ 1.4 μ l
X 35 μ l

0
0
X 28 μ l
X 1 μ l

4x23
= 92 μ l
RX B'113
MIX

ddw $n \times 16.15$
A'121 20X PCR $n \times 1.25 \mu$ l
B'89 4dNTP's 25mM $n \times 0.1 \mu$ l
B'89 DMSO $n \times 2.5 \mu$ l
14 μ l Left Primer $n \times 1 \mu$ l
14 μ l Right Primer $n \times 1 \mu$ l
B'113 Standard Tag $n \times 2 \mu$ l
B'113 E14 DNA 0.5 μ g/ μ l $n \times 1 \mu$ l
New T6-22 $n \times 25 \mu$ l
From Kim (8/4/94)

checked with Kim

or as suitable
500ng is
max. Can
use 100ng

can add later




Use $n = 14$

2 expect to set up
100 μ l RX for each primer set

Set up without primers.
Dispense 92 μ l to each of :-

	1L 1R B'113	1L 1MR B'113	1ML 1R B'113
B'113 RxMix	92 μ l ✓	92 μ l ✓	92 μ l ✓
Primer 1L	1L 4 μ l ✓	1L 4 μ l ✓	1R 4 μ l ✓
Primer 1R	1R 4 μ l ✓	1MR 4 μ l ✓	1ML 4 μ l ✓
Primer 1MR			
Primer 1ML			
Expected size	615 bp	432 bp	210 bp
Label	A	B	C

on 12:23 pm
35 cycles
Set up first at 93°
for ~ 12 min extra
Then 1 min 93°/92°
7 sec 40° dig
7 min 60° run
b.4.4 pm inspected
was finished Temp 46°
Shit bed at 5 pm

08286
ade: 8-4-94II: 23
: 52%08287
ade: 8-4-94II: 24
: 50%

Dye
W
● 1M L 1R
● 7L 8R
W
Acryl. MM
W
Dye
1.5% Hx1
On 10.27 am
Off 10.55 am
10V/cm

Primer
volumes
weren't

Conclusion: Something wrong
heating back - no check
marks on 20x PCR buffer!
But gel is not good
for 200bp. Re-run at 2%
O2 was new

Fri. Aug 5th

Intran 7 Clearance Receptor probe

- unknown length $\therefore$ use 10 min

PCR formula: -

~~PCR formula: -~~

Used the B'113 recipe for 14 tubes
~~dispensed to 93 μ l tubes~~

~~Divided into~~ $\frac{2}{3}$ & $\frac{1}{3}$

$\swarrow$ 207 μ l

$\equiv$ 9 tubes

+ 36 μ l **8R** ✓

+ 36 μ l **7h** ✓

$\swarrow$ etc to 8 $\times$ 35 μ l

Blue dot 

$\searrow$ Rest (115 μ l)

$\equiv$ 5 tubes

+ 20 μ l **IR** ✓

+ 20 μ l **IML** ✓

$\swarrow$ To 4 $\times$ 35 μ l

Black dot 

Started at $\sim$ 12:20 pm

93° for $\geq$ 2 min 1st cycle 1 min rest

40° 7 sec

60° 10 min

Expected finish

6:50 pm - out lab.

Note $\rightarrow$

Discarded
blue samples;
kept black



Conclusion: 1ML IR ^{probably} worked, but gel is not suitable
 Re-run (from B'115) at 2%.
 Put stained gel at 42° + stain overnight to look. (Confirms)

The blue product at 600bp is ~~very~~ conceivably correct.
 Would need to amplify with nested primers to tell.



Conclusion: 2% Seap. will separate well
 Don't know why B'115 is -ve.

Make new Seapaque

The Seap. was filtered thro'
 a ~~metal~~ metal
 filter - must have
 been contaminated.

just
 time
 but
 dug
 up
 a
 metal
 #7

Check for prep. gel

	Dye	
1ML IR	W	
1ML IR	Black	B'115
1ML IR	Black	B'117
	W	
	Marker	
	W	
	Dye	

1.5%
 Ag
 #7

2% Seap.
 HX
 75% Max
 (2.5 v/cm)

Sat. Aug. 6th

Repeat of 7L-8R CR

$n =$	14			
✓	226		ddw	16.15
✓	17.5	A'121	20X PCR	1.25
✓	1.4	B'89	4d NTP's 25mM	0.1
✓	35	B'89	DMSO	2.5
✓	28	B'117	Standard Tag	2
✓	14	B'113	DNA 0.5 μ g/ μ l	1
	322			23

Left primer | 1 |
Right primer | 1 |

✓	214 μ l (≈ 9.3 tubes)		107 (μ l) (Rest)
✓	9.3 μ l 8R		(≈ 4.7 tubes)
✓	9.3 μ l 7L		4.7 μ l 1R
			4.7 μ l 1ML

	Acc
	Water
15	1ML IR
17	7L 8R
	Water
	Acc. MM
	W

To $4 \times 55 \mu$ l To $2 \times 55 \mu$ l tubes

~ 1.05 pm started with ≥ 1 min extra melt
+ normal 1 min @ 93.5

Program 1 min 93
7 sec 40°


10:00 min 60°

Sh'd finish $\sim 7:35$ pm ✓

Cover
659.5°C

Dye
1.5 to HxI
Max 10V/cm
On 7.42 pm
Off 8.12 pm

Mon Aug 8th
 Prep. gel for IML1R



On 3:54 pm ≤ 15 V/cm Off ~4:40 pm

Only the B'117 had any band - but B'115 was a recovered sample anyway.


Stained. Cut out. Sylvie will recover

✓ It worked well.

B'117
 210bp
 Exon 1

B¹17 Exon 1 0.1KSSC

Panel
Bgl 2
EcoRI
Hind 3
Sca I
Xba I



434
marker

Mon. Aug 22nd

Two hypertensive papers off, back from the West Coast adventure, shingles 98% finished!

5' Clearance Receptor probe

S.H. tested B'117 fragment on Southern - reasonable with 0.1X SSC. But had better try to improve the chances. Work on rat L1 + mouse 1MR using different conditions &/or rat DNA.

Naomichi will test B'108 1 left with 1 Midnight with mouse & rat DNA using rather mild conditions: 94° 1 min

ON 11.25am off 1.5% H 10V/cm 50° 2 min 65° 5 min

+ DMSO

B'121 program

Dye	
Water	
21 Mouse DNA	12 1MR from Naomichi
22 Rat DNA	
10 7L 8R Mouse DNA	
Water	
AC MM	
Water AGMM	
Dye	

Conclusion: 7L/8R works well, as known. 1L/1MR maybe okay - see arrow - expected size is 434. Design next primer & try again.

le: 20
65%


5' note: 1L2 3'

C	T	A	T	T	G	C	C	C	C	G	G	G	A	C
G	A	T	T	C										

le: 20
65%

5' note: 1MR2 3'

C	C	C	A	T	C	T	T	G	G	C	G	T	A	G
G	C	A	G	G										



0.01 μ l
0.1 μ l
1 μ l

Tues. Aug 23rd

~~1L2~~ and ~~1MR2~~

Nested primers for 1L and 1MR.

Expected 383bp from the 434bp
using 1L and 1MR

Naomichi ran with B'121 program
→ got fine amplification - size slightly -
with 0.01µl, 0.1µl and 1µl of 1st round
1L/1MR amplification.
S.H. will make more a test.

Sequence: 08480

Total bases = 20

A= 3, G= 7, C= 6, T= 4, 5= 0, 6= 0, 7= 0, 8= 0
(mixed bases= 0)

MW: 6128.0

5'> CCC ATC TTG GCG TAG GCA GG <3'

1MR2

Sequence: 08479

Total bases = 20

A= 3, G= 5, C= 7, T= 5, 5= 0, 6= 0, 7= 0, 8= 0
(mixed bases= 0)

MW: 6065.0

5'> CTA TTG CCC CGG GAC GAT TC <3'


1L2

Mon. Aug 29th

Promoter Duplication

Over the last few weeks I have been considering how to compensate for $\downarrow$ in β globin expression consequent to selective for advantage genes within the locus. One possibility is to duplicate the whole gene ~~also~~ but cannot make a useful sickle cell model this way, or to put in an LCR or to duplicate the promoter.

Easiest test of this is with Neo (known to show effects of gene duplication -- higher [G418] resistance) can test in the HPRT locus in E cells: -



also available as HindIII

PMCI Neo^r portA

XhoI

CASSETTE

Enhancer (1)

HS

at

eu

II

2

TCGAGCAGTCTGGTTTTGCAAGAGGAAGCAAAAAGCCTCTCCACCCAGCCCTGCAATGTT
AGCTCGTCACACCAAAAACGTTCTCCTTCGTTTTTCGGAGACGTGGGTCCGGACOTTACAA

60

Enhancer (2)

TCCACCCAATGTGAGCAGTGTGGTTTTGCAAGAGGAAGCAAAAAGCCTCTCCACCCAGG
AGGTGGGTTACAGCTCGTCACACCAAAAACGTTCTCCTTCGTTTTTCGGAGAGGTGGGTCC

120

SV TK

Promoter

S
C
I
I
2

T
a
g
I
I
2

T
a
g
I
I
2

E
C
C
R
I
I
2

CCTGGAATGTTTCCACCCAATGTGAGCAAAACCCCGCCAGCGTCTTGTCAATTGGCGAAT
GCACOTTACAAAAGGTGGGTACAGCTCGTTTTGGGGCGGGTGGCAGAAACAGTAACCGOTTA

180

Red C See F101

N
S
P
V

M
S
I
I

TCCAAACACGCAGATGCAGTGGGGCGGGCGCGGTCCACAGGTCCATTGGCATATTAAGGTG
AGCTTGTGCGTGTACGTGAGCCCCCGCGCGCCAGGTCCAGGTGAAGCGTATAATTCCAC

240

A
f
IM
II
Iu
II

H
a
e
I

S
f
c
I

P
s
t
I

G
d
i
a
I
I

E
i
e
I
I

Neo

3-end

Bam
Sal

300

Hind

241

ACGCGTGTGGCCTCGAACACCGAGCGACCCTGCAGCCCAATATGGGATCGGCCATTGAACA
TGGGCACACCGGAGCTTGTGGCTCGCTGGGACGTGGTTATACCCTAGCCGGTAACCTTGT

Trans.
Initiation
Sequence

T
a
Gc


MBP

8

Thur. Sep 1st

Promoter & Neo Duplications


Thomas & Capra (Cell 51, 503, 1987) give
pMC1Neo as



Hind 3
or

Xho I - Pst I is the piece to duplicate 274 bp for
[Pro][Pro][Neo]

Xho I } Sal I or Hind I of pMC1Neo pA is the one
BamHI or




Cut with Xho I & insert Sal I into site. Check orientation (Sal goes into Xho)
by Xho cut plus Bam. $\overline{GTCGA}$ $\overline{CTCGA}$

Oligo # 2507Date made: 9-1-94LENGTH: 11% G+C: 64%

5'

C C T C G A G T G

note: OS PST → Xho

MW: 3315.2

5' > CCT CGA GTG CA <3'

Backup into
0.5 ml TE 1X C10A
Assume 1 µg/µl

CCTCGAGGTGCA

OS. Pst → Xho
B128



Uncl

Sal

Xho

Sal/Xho

Pst

Pst/Xho

on 5:41 pm
6:41 pm

0.8%
HX4
MAX

Dye

W

Uncl

129 Xho

129 Sal

129 Sal Xho

129 Pst

129 Pst Xho

W


Marker

W

Pyc


Conclusion:

Map must be: -



So cut with Pst, as planned, change Pst → Xho with
linker & go!

Check ΔHN for one (effectively) Xho site




Cut with Pst; expect to ↓ length by ~200bp
 - check against Xho cut. Change Pst to Xho.
 Excise Xho — Xho & put into ATIN

∴ take 33 µl (1.5 µg) ¹³⁰ ^{From NM (now labelled B'128)} ^{really} ^{0.46 µg/µl}
 MC1 Neo pol A SHI 114

✓ 14 µl 10x R1 salts
 ✓ 123 µl ddw

transferred to D'153
 moved to "Miscellaneous"



37° 2:35 pm → 5:25 pm

on 2:41 pm $\rightarrow 250 \mu A$
 $\rightarrow 5V/cm$ (total)
 off 3:42 pm

B¹³ Neo
sal/Xho
~40pb
1.8kb

190	0.8%
Scap	Scap Ag
1x	4x



5 } F
0 } M
5 } S

Error but -
proceed with

Neo/Sel/Xho

$\therefore + 80 \text{ p.e. } 5\text{M NH}_4\text{Ac}$
 Melted $\rightarrow + \phi\text{OH} / \text{HO}\phi$
 Then LXCIAA. stored as

B^{130} F	B^{130} M	B^{130} S
----------------	----------------	----------------

0.06
2009/12

B' (3)
KH2
Xho/C1AP
/C1AP

Whole of B131 KH21 Xho
for 10pe 1M Tris pH 8 (5x CIPF buffer)

✓ + 1 pe hab CIAP

30 min 55°C 3:37 pm 4:07 pm

1x HOQ/NOO
2x CIA

$$\frac{2x^2 + 2x + 1}{x^2 + 2x + 1}$$

Fri. Sep. 2nd

Neo/Neo & ProNeo (cont'd)

Take 11 μ l (5 μ g) B¹²⁹ pMCI Neo pol A ^{moved to D¹⁵³ 2 to "miscellaneous"}

5 μ l 10X R1 salts

34 μ l ddw

(~0.1 μ g/ μ l)

10 μ l (1 μ g)

+ ~0.5 μ l lot 39 Pst I
200/ μ l

B¹³¹
Neo/Pst

For linker ligⁿ

Rest (4 μ g)

+ 1 μ l Xho I lot 27 (200/ μ l)
+ 1 μ l Sal I lot 31 Pst (200/ μ l)

B¹³¹
Neo/Sal/Xho

For prep. gels
of Neo

Take 2.2 μ l (4 μ g) KH21 Φ 118^{2K} ^{Pro} Φ 118^{2K} $\rightarrow$ H (or vice versa)
H K

4 μ l Eco R1 salts

(~0.1 μ g/ μ l)

34 μ l ddw

B¹³¹
KH21
Xho

For CIAP

2.15 μ l (except Pst)

2 μ l (~0.2 μ g)

+ 2 μ l 10X R1 salts


+ 1/8 μ l
ddw

+ 1/2 μ l lot 46 (200/ μ l)
Hind 3

B¹³¹
KH21
Hind

For length
controls

2 μ l
Pst
KH21
Xho
to 32'



Conclusion:

linker-cond. appear
very low - but both
small bands got
larger.
B'130 M is visible.
B'131 KHE21 Xho bar
single cote.
Proceed.

On 11.5.7 am
off 1:12 pm
0.86 Hx4
MAR

	Dye	
	W	
+5	B'124 Pt	+5 pl A
2+16	B'133 Pt	linker
2+16	B'133 Pt	linker
Un dil. }	B'130 F	
	B'130 M	
	B'130 S	
Undil	B'131 KH	21 Hm
2+20	B'131 KH	Xho
	W	
	Martan	
	W	
	Dye	

4.5
no V/c
báo

Cho
m
a

Neo Neo ProProNeo (contd)

24 Pst + 10 pl At 1.04 pm

33 Pre linker ligation

33 Post linker lig

0 F

30 M

30 S

31 KH 21 Hind

31 KH 21 Xho

marker

ye

B'133
linker
ligation

Take whole (10 pl) of B'131 Neo Pst
+ 2 pl (1 μ g) B'128 OS Pst $\rightarrow$ Xho linker
+ 6 pl ddw
+ 2 pl p95 10x lig. buffer (2/3/86)
+ 1 pl T4 ligase

15° 1.14 pm to 1:54 pm

RT 1:54 pm to 2:15 pm

37° 2:15 pm to 2:50 pm

2 pl + 16 pl linker
Pre B'133 lig

2 pl + 16 pl linker
Post B'133 lig

Transformation 10 pl ($\sim 0.05 \mu$ g) to cold

80 pl ddw

10 pl 10x PBS

(Don't need purif. as the prep has $\sim$ zero ss DNA) $\rightarrow$ B'133 diluted

Transformation B'133 Neo Xho/Xho $\equiv$ Neo Δ XX

B'133 Dtt5 α 8/4 80 pl $\checkmark$
B'133 diluted 10 pl ($\sim 0.05 \mu$ g)

0° 3.12 pm to $\sim$ 3.50 pm

42°-30° 90 sec $\checkmark$

Plated Rest 20 5 20 5 37° 4 pm
(20) (9) (3) 16 16

Mems set up 1-25

To 37°

12:40 pm

B'133
Neo Δ XX

only just about enough

Processed to X vol but
also more for disc tests


Return to IX
Pen plated

Check
minis
w/ rest
Hind
and
Xho
at
B'133

1. Further
see much
12:40 pm

134


①

Mar
Sep 5th

← KH21

Conclusion: Pick some more! Run pre/post lig.
If none - restart. If 1 or more, pick more.

②



Conclusion:

It was not a
good lig.

Repeat with 200 µl B/30 M/
overnight in cold
Check prior to trans.

②

Dye

W

Pre 135

Post 135

W

Marker

W

Dye

Lig
lig

③



Conclusion:

lig. was
no good
by all
criteria

Top right B¹³⁵
repeated.

NO. 218
+ 16180.8
M5
OK

0.8% H₂O
 5:30pm Max 750mM
 off 6:25pm

Sat. Sep. 2nd

(1)

Dye

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288


289

290


291

292

293



Conclusion: Initial look suggests 2, 4, 6, 11 (12) 14, 19, 20, 24, 25
 1, 3, 7, 9, 10, 13, 15, 16, 17, 18, 21-23(?) prob. ~~short + Pst~~ are OK. - w'd be ~ 40% i.e. short + Pst
 Photo. again after RNA'se. Meanwhile
 digest 11, 14 & 12 with ~~thi~~ III & with Xho I



2 4 6 8 11 14 19 20 24 25 still look good.
~~12~~



Conclusion:

11 & 14 are okay.
 Can even use mini
 to make the
 Xho Euk Pro Xho
 fragment
 using 19 seap.

#14 streaked
 out as

B137
 PMCI
 ANeo

on 11.05 am
 off 12.07 pm
 0.8%
 HX
 max
 Dye
 w
 11 Hind
 11 Xho
 14 Hind
 14 Xho
 12 Hind
 12 Xho
 w
 Marker
 w
 Dye

in 5:05 pm
 of 5:50 pm
 0.8% HX4 750mM

Sun. Sep 4th

PK → Xho
 Xho

137

pmcI Neo Pat → Xho

Take

5 μ l (2x28) μ l 10x Eco RI

441 μ l ddw

1 μ l (0.25x28) (50/sample) hot 35 Pat I (200 μ l/ μ l)

25 x 18 μ l for [minis B'133 1-25]

32° at 12:32 pm

4:32 pm

B'137
 P1-25

B'137

Pat

1-25

minis.

6:30 pm

gel to ~55°
 with RNAse.

200 μ g + Tris buffer
 in ~400 μ l

B'137
 RNAse

6/12/94

10 μ g/ μ l

Take ✓ 2x8 = 16 μ l 10x Eco RI

✓ 16x8 = 128 μ l ddw

144

10 μ l

+ 1 μ l Xho I

hot 27

18 μ l

Rest

+ 1 μ l Hind

hot

18 μ l

+2 μ l

min

11

14

12

X

X

X

+2 μ l

min

11

14

12

H

H

H

32° 6:55 pm to ~9 pm


19 μ l B'133 diluted
 Act Marker
 W
 Dye

25
 Part - Pat

5 am
 07 pm
 0.8%
 HX4
 max

Hind
 Xho
 Hind
 Xho
 Hind
 Xho


marker
 N
 Dye



Conclusion:

Difficult to believe -
I have run with
B¹³³ del. control.

Cut out a 2.2 g
band anyway
at ~2.3-2.8 cm - frozen
Prob. but ~50%
Prepare next
- later.



Conclusion: The digests of
pooled minis worked fine.
Prep gel did not. Use 0.8%
Ag Hx4 & cut for electroelution.

The B¹³⁵ digⁿ worked well
- but (as expected) a lot
of circularization of insert.
If unsuccessful, repeat at
higher concⁿ. Meanwhile
transform & screen.
Also ↓ amount of KH21.

← Same mixed
Reversed

On
of
8+8
8+8
Undil
4+10
4+1

Tues. Sep 6th

Prep. gel for Xho Euh. Pro Xho

- Take 40 μ l 10x RI salts
 ✓ 280 μ l ddw
 ✓ 40 ~~280~~ μ l of B¹³³ mini #25
 ✓ 40 μ l of B¹³³ mini #11
 400 μ l


✓ +8 μ l (160v) ✓✓ 8+8 Pre ~~280~~ Xho

Xho Lot 27

32° 3:46 PM to 5 PM

8+8 Post Xho

B¹³⁹
Xho



July 12, 1992 18:52

Also available as Hind 3

Xho I

May need hot start to avoid this priming but only 2 or 10 Bp.

HS
at
eu
II
/

T
a
q
I
I
-
2

Not Y

TCGAGCAGTGTGGTTTTGCAAGAGGAAGCAAAAAGCCTCTCCACCCAGGCCTGGAATGTT
AGCTCGTCACACCAAAAACGTTCTCCTTCGTTTTTCGGAGAGGTGGGTCCGGACCTTACAA

FORWARD

ENH

TCCACCCAATGTCGAGCAGTGTGGTTTTGCAAGAGGAAGCAAAAAGCCTCTCCACCCAGG
AGGTGGGTACAGCTCGTCACACCAAAAACGTTCTCCTTCGTTTTTCGGAGAGGTGGGTCC

S
C
U
I
/

T
a
q
I
I
-

T
a
q
I
I
-

E
C
C
O
R
I
I
/

CCTGGAATGTTTCCACCCAATGTCGAGCAAACCCCGCCAGCGTCTTGTCAATTGGCGAAT
GGACCTTACAAAGGTGGGTACAGCTCGTTTGGGGCGGGTCGCAGAACAGTAACCGCTTA

N
S
P
V

M
S
I
I

TGGAACACGCAGATGCAGTCGGGGCGGGCGCGGTCCAGGTCCACTTCGCATATTAAGGTG
AGCTTGTGCGTCTACGTCAGCCCCGCGCGCCAGGGTCCAGGTGAAGCGTATAATGCCAC

A
f
IM
II
IU
II

H
e
e
I

S
f
a
I

P
S
C
I

G
d
i
I
I

Reverse


Met

ACGCGTGTGGCCTCGAACACCGAGCGACCTGCGGCCAATATCGGATCGGCCATTGAACA
TGCGCACACCGAGCTTGTGCTCGCTGGGACGTCGGTTATACCCTAGCCGGTAAGTTGT

Pro

25 No
Cater!

Tues. Sep. 6th
Diagnosis of KH21Neo^R Enh^R Pro^R



Inversion 1 primer will work
Direct Repeat Needs 2 primers
Can check on B137 (pMC1) ΔNeo Pet
Back into 0.5ml TE/CIAA assume 1pg/μl

B141
FOR ENH

B141
REV PRO

5'	Oligo Name:	FORWARD ENH B141	3'
le:		T T T C C A A A T G T C G A G C A G T	
li:	22 50%	G A G	
		8568	
5'	Oligo Name:	REVERSE PRO B141 (G T C)	3'
le:		G C C A	
li:	22 50%	T T A A	
		8568	

5' > TTT CCA CCG AAT GTC GAG CAG T <3'

5' > GCG ACA CGC GTC ACC TTA ATA T <3'

HAT	1X G418	1X G418 +	1.7X G418	1.7X G418 +	3X G418	3X G418 +
		HAT		HAT		HAT

KH21	15	1170	13	1230	24	1060	14
	14	1170	11	1070	19	820	14

NeoDnp	1660	20	1210	18	650	20
	1400	14	1240	14	840	19

2PNeo	1390	17	1390	22	850	16
	1150	20	1050	18	740	17

SH U78 DNAs were:—

KH21 ϕ 118²
NeoDnp U75 #92

2PNeo SH U64 #78 (B'147 $\rightarrow$ #78 U47)

12/29/94

Conclusion: This assay detects no differences. $\therefore$ picked 10-20 of HAT^R G418(1x)^R $\rightarrow$ some G418^R for use in assay of Neo activity.

Wed. Sep. 7th

Neo Neo (repeat)

Transformation



B'143 DH5 α 8/4	80 μ l
B'135 2 lig ⁿ	4 μ l
0°C	12:15 PM to 1:08 PM
90 sec. 42°	2 min
Plated	on (1X) Amp NZY (Rest) 20, 5, 20, 5 16, 16
	on (2X) Amp 20 37° 1.20 AM to 1:00 AM

Precautionary repeat ligⁿ. : — Thurs. Sep 8th

✓ 2 μ l (i.e. 25%) B'131 KH₂1 K_{ho}/CIA θ /CIAA
 ✓ 200 μ l B'130 M
 ✓ 22 μ l 8 M NH₄Ac
 ✓ 672 μ l EtOH
 Dry Ice, Cent. Dry.
 i.e. 667 — Back into ~~90 sec~~ 10 μ l 1X p95 ligⁿ. buffer
 1 μ l T4 ligase B'135
 Cdd 9:35 am to 4 pm Fri
 B'143 KH₂1 Neo ligⁿ
 2+14
 At 143 ligⁿ
 2+14
 Post 143 ligⁿ

See SH U75 for Neo Neo KH₂1 product.

Restriction map of mouse angiotensinII receptor (Type 1A) gene




Tues. Nov. 15th Grants & papers & finished!

ATRIA Duplication

Masaki & Tom's success with ATRIA disruption
 & its effects on bp. (↓ ~~mut~~ Hg in +(-) → lack
 of ATRIA responsiveness in -/- suggests that dupⁿ
 will ↑ bp. & lost same concepts as ↑ ANPR-CR
 clearance receptor
 (see B'107)


Can use the same 3' piece as Masaki -
 except for putting in a diagnostic site that will work
 with Southern for 2/1 versus 2/2. In fact no site
 within the duplication will work except quantitatively.
 Once again, either a 129 v. B6 difference is needed
 or duplication must be within scale of outside
 sites. Could use the Bam sites as Masaki did & duplicate



Can also look for 129 v. B6 polymorphism.

(In)compatibility of ends


Eag/Not



Xba




Bam/Bgl





Sal

All four
are discrete.

Wed. Nov. 16thInitial plan for constructlinearize with Xho

Homology ~ 8 kb.



← Primer for PCR detection can be made from within

Eag H Xba

→ Primer w'd be from 3' Neo.

PCR frag^t ~ 2.5-3 kb.Comment

The 5' duplicated gene will be very secure. The 3' copy may be somewhat short in promoter end.

5' Oligo Name: OSXBABAM 3'

C	T	A	G	C	G	G	A	T	C	C	G			

12
67%

5' Oligo Name: OSBAMSAL 3'

G	A	T	C	G	G	T	C	G	A	C	C			

Cancelled

12
67%

5' Oligo Name: OSXHOBGL 3'

T	C	G	A	C	G	A	G	A	T	C	T	C	G	

14
57%


Into 0.5 ml TE
1x CIAA
→ labelled

Thurs. Nov. 17th

Same adapters


XbaI → BamHI

OSXBABAM B'149




Bam → Sal

OSBAMSAL B'149



Xho → Bgl2

OSXHOBGL B'149



Transformation

✓ DH 5 α 8/4 B'143	80 μ l
✓ B'151 LIG ^N	20 μ l $\cong$ 0.1 μ g
✓ Ice 3:15 pm to 3:48 pm	
✓ 43°-42°	90 sec
plated	Rest 20 5 $\frac{20}{16}$ $\frac{5}{16}$
2X Amp	
37° 4:02 pm	B'151 Neo Bam


Picked 12 (with difficulty)

37° at 8:45 am


N.M. already had made pMCINeo polA
without Bam, & it cuts out with Xho/Sal

Thurs. Nov 17th


pmc1Neo Bam



Can simplify process without linker by cut & fill.



- + 1 µl 1mM mix of dNTP's B'151 4 dNTP's 1mM each
- + 1 µl Klenow polymerase 30 µl lot 57



Gel → over



Conclusion:

lighting & film were good.

Pool 0, 1, 2, 4, 8

& prepare single

cut, or repeat

in bulk at 2 min.

On 3:40 pm
Off 4:55 pm
0.8% H₂O max

Dye
W

Pre
"O"

1

2

4

8

16

32

R

W

Marker

W

Post 2 lig^m


Pre B'151

Post 2 Bam

Pre B'151

W

Dye



On 11 35 am
400 mA Off 4:50 pm
0.3 V/cm

Sample is whole of B'153 PI-E7 X → B
+100 µl ddw ✓

Dyes

B'152

1400
1200
1000

F
M
S


Washed

+ 10th 8M NH₄Ac

Melt 85°

1 x POH/HOQ

2 x CIPA



1x4
max↓ to 2/3
at 4pm↑ 1x
4:30pmThurs. Nov 17th

Xba



Xba → Bam

Partial digests of P1-E7

0.05 pg/μl
 B'153
 Xba I

Take 5 μl (~10 pg) P1-E7 (1) 2.1 pg/μl 2/14/92
 + 20 μl 10xRS + 1.5M NaCl
 + 175 μl ddw
 + 1/2 μl (10u) Xba I hot

← to help with heat inact
 20 μl

~(1:55pm)

Pre	0	1 min	2 min	4 min	8 min	16 min	32 min	57 min
20 μl	✓	✓	✓	✓	✓	✓	✓	Rest
To 85° for 5 min.	90°							

Each time point 1 μl, after heating, 2 μl + 16 μl for exp.

Pooled 0, 1, 2, 4, 8 to total vol. of ~75 μl

75 μl Partial Xba P1-E7 (~5 pg)
 49 μl 10xRS 95 ligase buffer
 + 5 μl B'148 0.5 XBA BAM
 + 2 μl T4 ligase

15° 255pm to 3.55pm
 RT 2.40pm to 2.55pm 37° 3.55pm to 4.4pm
 Träfen

Use now for prep. gel



Conclusion:

NM₂ material
is as advertised!

Abandon B'150/1510
expt. & proceed
with B'155

On 2:15 pm

~~Off~~ 4:10

Dye

W

Pre

Post X

Post X

Post X

W

Marker

W

Dye

1+15

B'154

#3

straked out (see B'156)

ad

B'154

B'Neo S

80 pl B'133

CA hole of
B'155 Neo Bgl
Heat

0° 5.15 to 5.45 pm

42°

90 sec

Plated

Rest 20, 5, 20, 5
16 16

37° at

#2

#1

#3-5

3 v. small

1

1

OK

#1-5 B'154

37° 10:20 am

(negative)

As control for Bgl 2

37° 11 am, 11:42 am

took B'155 X+B & added 1/2 pl Bgl 2

1+16


Pl B'154
Bgl 2

Post B'154
Bgl 2

5pm
4:10

0.8% various
HX4 voltages

Fri. Nov 18th



155

mg p
re
Xho
Xho
Xho
Marker
Dye

B'155
+ Bam
+ Sal

Xho pMC1Neo Bam
↓
Bgl2

Take 4 μ l ($\sim 2 \mu$ g) X135 #1 Midi (B'155)
(pMC1Neo posA minus Bam
from N.M.)

B'155
Neo
Xho

+ 4 μ l 10x R1 salt
+ 32 μ l ddw

20 μ l + $\frac{1}{2}$ μ l (100) Xho I lot

10 μ l + $\frac{1}{2}$ μ l Bam^{HI} lot


1+15

Pre
B'155

37° at 9:10am to 11:25am

10 μ l + $\frac{1}{2}$ μ l (100)
1+15 Sal I

Post B'155
X+S



1+15
Post
B'155
X+B

whole
Take 19 μ l ($\sim 0.6 \mu$ g) pMC1Neo Xho

0.03 μ g / μ l
B'155
Neo
→ Bgl2

+ 2.4 μ l 10x PBS high buffer
+ 1 μ l ($\sim 1 \mu$ g) B'148 OS XHO BGL
+ 1 μ l T4 ligase

RT 2:40pm to 2:55pm

15° 2:55pm to 3:55pm 37° to 4:45pm


3.3 μ l + 2 μ l 10x PBS
0.1 μ g + 15 μ l ddw

85° for 3 min

B'155
Neo
Bgl
Heat

B'154

Conclusion: #3 mini has Age 2 - no site
in original plasmid // Fragment M
is good for ~~Xba~~ — Xba
↓
Bam HI



Sun. Nov 20th

Minis etc.

Processed B'154 mini#1-5.

Take ✓ 12 pl 10X R1 salts

✓ 107 pe d d w

✓ 1 pl (200) Bgl 2 hot

B'157
B 1-5

1 pl of each min to 20 pl 37° 12.15 pm to 2:53 pm
not fully dissolved

B'152

F 280 pl + 0.84 ml

M 240 pl + 0.72 ml EtOH

S 200 pl + 0.60 ml

10% F into 20 pl
10% M
10% S !

on 3:50 pm
5:15 pm

0.8% H x 4 max

Aye
W

Post Bgl 2 B'154 (Xho + Bam)

Post

1

2

3

4

5

(B'154 Post Bgl 2)

mini

B'157 #1-#5

Load B'152 prep gel

10% F

10% M

10% S

W

marker

W

8yl

①



processed minis as usual ~ 10am.
 I digest is simplest initial screen for absence
 of Xba I site. Bam is better for +ve, but
 know if the vector has a site.

54 μ l 10x R~~S~~ + 1.5M NaCl
 + 480 μ l ddw
 + 6 μ l (120 u) Xba I hot 29 (20u/ μ l)
 to 20 μ l. Each + 1 μ l minis ^{Pipetting was a little erratic.}

59 #1 - #25 37° 11:35am to 2:15pm

① Conclusion: 60% successful! Pick ~5 for
 orientation.

#s 1, 2, 3, 6, 7 & 12
~~will do fine.~~

(Error)

$\therefore$ take 14 μ l 10x RS + 1.5M NaCl

✓ + 126 μ l ddw

✓ + 1 μ l (20u) hot 29 Xba I


✓ + 43 μ l Sal I

B'159

minis 1, 2, 3, 6, 7 & 12

See B'162
 for finals

②



② Conclusion: #3 and #12
 can go on to full test

②
 Dye
 Water
 1
 2
 3
 6
 7
 12
 W
 W
 W
 Dye

B'159
 Xba
 Sal
 min

On 3pm
Off 4.05pm
0.86 Hx4 Max

Sam. Nov 20th

Euker → Ban
~~Xba~~ → ~~Xba~~

or → Ban
~~Xba~~

Dye (1)

~10 pl (equal to gel load) of B'157 "10% M" available
i.e. $\approx \frac{1}{20}$ th.

Added 1 pl 10xPBS, heated to 85° for 3 min, iced
→ to 80 pl bacteria.

Also 80 pl of crude B'152 M heated to 85° for 3 min, iced
→ to 80 pl bacteria

	Conc?	Grade
DH5α B'133	80 pl	80 pl
DNA	11 pl "10% M" B'157 heated	80 pl B'152 M (crude) heated
Lce	6.22 pm	6.53 pm
42°	90 sec	

Paired Rest, 20, 5, 20, 5

or 1x amp

as B'159 Crude

B'159 conc.

9:15 am Mon. Zero on "Crude"
Excellent on "10% M"

Minus B'159 #1-25 from ~10:30 am to 7:30 pm
with shaking. R.T. from then on.

Bgl 2

Hx4
max
6.45 pm
1.45 pm

(2)

Dye
Water

1

2

3

6

7

12

W


M

W

Dye

Marker
Dye

B'159
Xba
Sal
min




Sol Kba Pam Sol Kba Pam Sol Kba
 3 3 3 12 12 12 12 12 12 12 12 12

on 11.38 am
12.54 pm
10.86 HX
max.


(1)

Dye
w
#3 Sal
#3 Xba
#3 Bam
#12 Sal
#12 Xba
#12 Bam
#3 } Sal
#12 } Xba
#3 } Sal
} Bam
7 Xba
Bam

SacI SacI
3 12



3	12	3	12	3	12
Sact	Sact	Sac/Kda	Sac/Kda	Sol/Kda	Sol/Kda



try orientation again with Bgl 2

$$\begin{array}{r} \text{Dye} \\ \hline \text{W} \\ \hline \#3 \} \text{Sac} \quad \text{I} \\ \#12 \} \\ \hline \#3 \} \text{Sac} \quad \text{I} \\ \#12 \} \text{W} \quad \text{I} \\ \hline \#3 \text{ Sol} \\ \#12 \times 68 \quad \text{I} \\ \hline \text{W} \\ \text{W} \\ \hline \text{W} \\ \hline \text{Dye} \end{array}$$

↓
1pt #3
Sal
nbw

0.89M
max

I

$\frac{I}{I}$

↓ Dist
rest
of

+14


Wed. Nov 23rd
Checks on Xba

Bam

(161)

B'159 #3 & #12
minus

M1 Take 30 ~~pl~~ 10x RS+15M ~~pl~~
✓ 270 ~~pl~~ 2 dw



Error - I have used Sac I for orientation

37° 10:25am to 11:25am

B'159
#3

B'159
#12

streaked out at 37° Norm Wed.

Need to recheck with Xba I and Sac I for orientation;
Sal I was probably in the vector.

Take 10pl R1 salts
+ 90pl ddw
+ 1pl ~~Sac I~~ Lot 45

50pl
+ 1pl ~~Xba I~~ Lot 29

20pl
+ 1pl #3 Sac

#12 Sac

#3 Xba Sac

#12 Xba Sac


37° 12:10pm
21:30pm

162



Thanksgiving Day, stained & photographed.

Conclusion: $\leftarrow 2 \rightarrow \leftarrow 3.5 \rightarrow$




Final maps are: —


#3 B'159



This was the one planned. Can use for



#12 B'159



Can use for Bam — Xba
Bam — Sal



ad 8
graphed.

5 →

Bam
Xba

Xba

Xba1

BamH1

mm

on 7:05 pm
off w 8 pm
0.86 #x4
max

Wed - Nov. 22nd, 1994

Orientation of #3 & #12 B'159 (contd.)

Take ✓ 10pe 10xR1 salts
✓ 90pe 1dw
✓ 1pe 10xII lot 34

Rest
+0.6
XbaI lot 29

Dye
w
#3 } Bgl2
#12 }
#3 } Bgl2
#12 } XbaI
w
marker
w
pge

20 20
#3 Bgl #12 Bgl #3 B/x #12 B/x
37° 6pm

Boorum & PeaseTM

Columnar Books • 10 3/8 in. x 8 3/8 in.

150 Pages	300 Pages	Ruling	Width of Desc. Space	Units Per Column
Record Ruled				
21-150-R	21-300-R	Record	-	-
Columnar / Ledger / Quad Ruled				
Single Page Forms				
21-150-2	21-300-2	2 Columns	3 3/8"	10,000,000
21-150-3	21-300-3	3 Columns	3 3/8"	1,000,000
21-150-4	21-300-4	4 Columns	2 3/8"	1,000,000
21-150-5	21-300-5	5 Columns	1 5/8"	100,000
21-150-6	21-300-6	6 Columns	1 3/8"	10,000
21-150-7	-	7 Columns	1 3/8"	10,000
21-150-DEL	-	D.E.Ledger	1 3/8"	10,000,000
21-150-Q	21-300-Q	Quad-5 Sq.	-	-
Double Page Forms				
21-150-8	21-300-8	8 Columns	3 5/8"	10,000,000
21-150-10	21-300-10	10 Columns	4 1/4"	1,000,000
21-150-12	21-300-12	12 Columns	2 3/8"	1,000,000

ESSELTE
Esselte Pendaflex Corporation, Garden City, NY 11530 • Made in U.S.A.